



# Abundance of Major Insect Pests in Rice and Their Relation to Weather Parameters

Arvind Kumar <sup>a\*</sup>, Pankaj Kumar <sup>a</sup>, Umesh Chandra <sup>a</sup>,  
Pradip Kumar Patel <sup>a</sup>, Vishnoo Omar <sup>a</sup>,  
Ratnakar Pathak <sup>a</sup> and Ravi Kumar Rajak <sup>a</sup>

<sup>a</sup> Department of Entomology, College of Agriculture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya – 224229, Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i82129

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100886>

**Original Research Article**

**Received: 25/03/2023**

**Accepted: 31/05/2023**

**Published: 16/06/2023**

## ABSTRACT

The present investigation has been conducted in rice crop during *Kharif* 2021-22 at Students' Instructional Farm of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.), India. The population of six major insect pests was recorded *viz.* Rice Yellow stem borer (*Scirpophaga incertulus*), Rice Leaf folder (*Cnaphalocrocis medinalis*), Rice ear head bug (*Leptocorisa acuta*), Brown plant hopper (*Nilaparvata lugens*), White backed plant hopper (*Sogatella furcifera*) and Green leaf hopper (*Nephotettix nigropictus*) that cause most damage at different crop growth stages initiated after sowing till harvesting. The maximum population of yellow stem borer was observed on rice crop at 36<sup>th</sup> SMW (12.20 per cent dead heart). The infestation of dead heart was non-significant positively correlated with minimum, maximum temperature and RH% (0.433, 0.010 & 0.253 respectively) and negatively correlated with rain fall (0.081) whereas,

\*Corresponding author: E-mail: ak847051@gmail.com;

significant negatively correlated with sunshine (0.552\*). The maximum population white ear in 40<sup>th</sup> SMW (16.80 per cent). The infestation of white ear was non-significant negative correlation with minimum, maximum temperature (-0.062, -0.167) and positive correlation with RH & rain fall (0.244 & 0.173) whereas, significant positive correlation with sunshine (0.703\*). Maximum leaf folder population was observed during 35<sup>th</sup> SMW (16.35 per cent). and infestation of leaf folder was non-significant positive correlation with minimum temperature & RH (0.427 & 0.505) and negative correlation with maximum temperature, rainfall and sunshine (-0.079, -0.030 & -0.291). The maximum population was recorded of ear head bug in 38<sup>th</sup> SMW i.e., (10.90 ear head bug/hill respectively.). The infestation of ear head bug was non-significant positive correlation with minimum temperature, RH, rainfall & sunshine (0.061, 0.176, 0.155 & 0.210) and negative correlation with maximum temperature (-0.407). The peak mean population of BPH was recorded in 37<sup>th</sup> SMW i.e., (11.25 brown plant hopper/hill respectively.) It was non-significant positive correlation with minimum temperature & RH (0.463 & 0.357) and negative correlation with maximum temperature & rainfall (-0.066 & -0.088) whereas, significant negative correlation with sunshine (-0.597\*), The peak mean population of WBPH was recorded in 38<sup>th</sup> SMW i.e., (12.80 white backed plant hopper/hill respectively.) It was non-significant positive correlation with minimum temperature, RH & rainfall (0.300, 0.337 & 0.185) and negative correlation with maximum temperature (-0.300) whereas, significant negative correlation with sunshine (-0.094\*) and the maximum population of green leaf hopper was recorded in 37<sup>th</sup> SMW i.e., (10.80 green leaf hopper/hill respectively.) with the infestation of green leaf hopper was non-significant positive correlation with minimum temperature & RH (0.479 & 0.369) and negative correlation with maximum temperature & rainfall (-0.077 & -0.120) whereas, significant negative correlation with sunshine (-0.633\*).

*Keywords: Rice; major; insect pests; weather parameters.*

## 1. INTRODUCTION

“Rice (*Oryza sativa*), is the most important crop of the world which occupies foremost status in human food requirement. It is the staple food of more than 60 per cent of the world’s population because of the major source of calories. About 90 per cent of all rice grown in the world is produce and consumed in the Asian region. Rice crop is grown worldwide over acreage of around 167.25 million hectares” [1]. “It is cultivated in almost all tropical, subtropical and temperate countries of the world. Almost 90% of rice is produced and consumed in Asian countries like China, India, Japan, Korea Republic, Sri Lanka, Pakistan, Bangladesh, etc”. (Nadaf et al., 2016). “The major rice growing countries are India, China, Japan, Indonesia, Thailand, Burma, Philippines and Bangladesh. India is amongst the top most rice producers in the world, second only to China. Rice contributes more than 40 per cent of the country's total food grain production” (Anonymous, 2018). “According to the data released by the Government of India, the annual production of rice in India is 115.63 million ton during 2018-19 [2]. “The protein content of milled rice is usually 6 to 7 per cent. Rice however”, compares favorably with other cereals in amino acids content. The biological value of its protein is high. The fat content of rice is low (2.0 to 2.5 per cent) and much of the fat is lost during

milling. Rice contains a low percentage of calcium. The byproducts of rice milling are used for various purposes. Rice bran is used as cattle and poultry feed. Rice husk can be used in manufacture of insulation materials, cement, card board and as a litter in poultry keeping. Rice straw can be used as cattle feed as well as litter during winter. Rice yield is affected by many factors in which, climatic conditions and pest epidemics are most important. Reduction in the rice yield is due to many biotic and abiotic stresses such as, pests, diseases, soil fertility, rainfall, water logging and climatic conditions. Amongst the various threats to rice productivity, damage by various types of insect pests is considered significant as it can cause large scale reduction both in quality and quantity of rice. Such cultivation pattern of rice accidentally or in advertently offers infestation of a large number of insect pests, which results in to severe loss in crop yields” [3]. “Weather plays an important role for determining the geographical distribution and periodic abundance of major insect pests in rice” [4]. Among the weather factors; temperature, rainfall, relative humidity plays the crucial role in insect life. Hence, considering the importance of insect pests of rice an attempt has been made to study the influence of different abiotic factors on the incidence of major insect pests of rice.

## 2. MATERIALS AND METHODS

The present investigation was conducted under field conditions at students' Instructional Farm, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, Uttar Pradesh, India during *Kharif*, 2021-2 on rice variety of NDR-2065. The quantification of the insect pests population was done under field conditions in the plot size kept 4x3 meter with border 1 meter and line to line spacing kept 20 cm and plant to plant spacing kept 15 cm. The details of experiment are given in. The occurrence of the insect pests was recorded 20 days after transplanting (DAT). The pest population were recorded randomly selected 10 hills per plot at weekly intervals to pre-harvesting. The meteorological data of weather parameters were obtained from the Department of Agricultural Meteorology, College of Agriculture, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, (U.P.). The correlation coefficient between insect pests population and weather parameters like maximum and minimum temperature (°C), relative humidity (%), rainfall (mm) and sunshine (hrs.) was worked out.

## 3. RESULTS AND DISCUSSION

### 3.1 Yellow Stem Borer (*Scirpophaga incertulas*)

Yellow stem borer was recorded at weekly intervals during crop period revealed that (*S. incertulas*) appeared at vegetative stage crop is called dead heart. During *Kharif* season, 2021, the initial infestation of (*S. incertulas*) was recorded in 29<sup>th</sup> SMW (0.50 per cent dead heart) and increase up to peak level of (12.20 per cent dead heart) in 36<sup>th</sup> SMW. Thereafter, yellow stem borer of declined and it was not appeared 41<sup>th</sup> SMW. It was non-significant positively correlated with minimum, maximum temperature & RH (0.433, 0.010 & 0.253 respectively) and negatively correlated with rain fall (0.081) whereas, significant negatively correlated with sunshine (0.552\*). Yellow stem borer was recorded at weekly intervals during crop period revealed that (*S. incertulas*) appeared at panicle initiation stage of crop is called white ear. During *Kharif* season, 2021, the initial infestation of (*S. incertulas*) was recorded in 36<sup>th</sup> SMW (0.70 per cent white ear) and increase up to peak level of (16.80 per cent white ear) in 40<sup>th</sup> SMW. Thereafter, yellow stem borer of declined and it

was not appeared 43<sup>th</sup> SMW. The infestation of white ear was non-significant negative correlation with minimum, maximum temperature (-0.062, -0.167) and positive correlation with RH & rain fall (0.244 & 0.173) whereas, significant positive correlation with sunshine (0.703\*), (Table 2 and Fig. 1). These findings are similar with the finding of Jasrotia et al. [5] notice that maximum damage of 14.9% and 14.5% by yellow stem borer was recorded during 38<sup>th</sup> SMW in 2016 and during 39<sup>th</sup> SMW in 2017, respectively. These finding also similar to Patel and Singh [6] reported that incidence of rice stem borer on different stage of crop. Dead heart started from 32<sup>th</sup> standard week and continued up to 38<sup>th</sup> standard week, while white ear head was recorded on 35<sup>th</sup> standard week and it reached peak in 45<sup>th</sup> standard week.

### 3.2 Leaf Folder (*Cnaphalocrocis medinalis*)

The leaf folder infestation was noticed in different crop periods. During *Kharif* season, 2021, the initial infestation of (*C. medinalis*) was recorded in 29<sup>th</sup> SMW (1.10 per cent) and increase up to peak level of (16.35 per cent) in 35<sup>th</sup> SMW. The infestation of leaf folder was non-significant positive correlation with minimum temperature & RH (0.427 & 0.505) and negative correlation with maximum temperature, rainfall and sunshine (-0.079, -0.030 & -0.291), (Table 2 and Fig. 1). These findings are partial agreement with the finding of Jasrotia et al. [5] notice that infestation of leaf folder attained peak during 3<sup>rd</sup> week of September (38<sup>th</sup> SMW) during 2016 and 2017 with 8.4% and 9.7% infestation levels, respectively. These finding are also similar to Priya and Prasad [7] record the incidence of leaf folder started during 34<sup>th</sup> standard week with 13.0% leaf damage. Then, the infestation increased gradually and attained its peak in 40<sup>th</sup> standard week with 21.4% leaf damage.

### 3.3 Ear Head Bug (*Leptocorisa acuta*)

The ear head bug population was observed at panicle initiation & milking stage of crops. During *Kharif*, 2021, the initial population of (*L. acuta*) was recorded in 35<sup>th</sup> SMW (1.95 bugs per hill) and increase up to peak level of (10.90 bugs per hill) in 38<sup>th</sup> SMW. The infestation of ear head bug was non-significant positive correlation with minimum temperature, RH, rainfall & sunshine (0.061, 0.176, 0.155 & 0.210) and negative correlation with maximum temperature (-0.407), (Table 2 and Fig. 1).

**Table 1. Occurrence of major insect pests on Rice in relation to weather parameters during Kharif 2021-22**

SMW	Incidence of Major Insect Pests							Weather Parameters				
	Yellow stem borer		Leaf folder (%)	Ear head bug/hill	BPH/hill	WBPH/Hill	GLH/hill	Temperature (°C)		Average RH (%)	Rainfall (mm)	Sunshine (hrs.)
	Dead heart (%)	White ear (%)						Min.	Max.			
29.	0.50	0.00	1.10	0.00	0.20	0.00	0.00	21.80	34.10	80.00	38.00	5.30
30.	0.80	0.00	1.30	0.00	0.80	0.50	1.20	27.20	34.40	79.60	37.40	5.60
31.	1.00	0.00	2.50	0.00	1.45	0.90	4.50	26.40	33.20	76.00	30.00	5.90
32.	2.15	0.00	5.80	0.00	3.70	1.50	8.60	26.20	32.20	85.90	12.50	3.40
33.	4.70	0.00	8.60	0.00	5.20	1.80	10.20	26.20	38.80	80.80	18.80	5.40
34.	6.25	0.00	13.05	0.00	8.75	2.35	17.60	25.50	32.20	87.80	24.60	2.60
35.	9.32	0.00	16.35	1.95	9.20	5.30	18.00	25.80	32.70	81.20	1.00	5.50
36.	12.20	0.70	11.20	5.20	10.60	6.70	19.20	26.20	33.70	79.40	24.20	3.90
37.	8.25	1.00	11.80	10.00	11.25	10.35	20.80	25.50	32.10	78.80	2.60	4.40
38.	6.50	3.80	9.90	10.90	7.45	12.80	16.30	25.20	31.20	85.80	41.40	6.10
39.	3.30	9.70	12.40	7.25	4.50	7.30	6.20	25.10	32.20	84.50	63.00	7.30
40.	0.50	16.80	8.40	4.50	1.80	3.70	2.25	24.40	32.70	82.80	18.00	8.10
41.	0.00	7.50	4.60	3.50	0.00	1.33	0.20	24.00	34.00	79.50	0.00	9.00
42.	0.00	4.20	2.20	2.80	0.00	0.00	0.00	23.60	31.80	81.30	24.00	6.90
43.	0.00	0.00	1.31	1.50	0.00	0.00	0.00	18.20	31.10	68.00	0.00	6.50

SMW= Standard Meteorological Week




Fig. 1. Occurrence of major insect pests on Rice in relation to weather parameters during *Kharif* 2021-22

**Table 2. Relationship between insect pests of Rice and weather parameter during Kharif 2021-22**

Insect Pests		Weather Parameters				
		Temperature( C)		Relative Humidity (%)	Rainfall (mm)	Sunshine (hrs.)
		Min.	Max.			
Yellow stem borer	%DH	NS (0.433)	NS (0.010)	NS (0.253)	NS (-0.081)	-0.552*
	%WE	NS (-0.062)	NS (-0.167)	NS (0.244)	NS (0.173)	0.703*
Leaf folder		NS (0.427)	NS (-0.079)	NS (0.505)	NS (-0.030)	NS (-0.291)
Earhead Bug		NS (0.061)	NS (-0.407)	NS (0.176)	NS (0.155)	NS (0.210)
BPH		NS (0.463)	NS (-0.066)	NS (0.357)	NS (-0.088)	-0.597*
WBPH		NS (0.300)	NS (-0.300)	NS (0.337)	NS (0.185)	NS (-0.094)
GLH		NS (0.479)	NS (-0.077)	NS (0.369)	NS (-0.120)	-0.633*

\*Significant at 5%

These findings are in agreement with Girish et al. [8] found that population gradually increased of earhead bug on 38<sup>th</sup> standard week as 4 nos./10 sweep nets. Later, the population suddenly increased with 9 nos./10 sweep nets when the panicle emerged and gradually reached to 17 nos./10 sweep nets during 44<sup>th</sup> standard week. These findings also similar to Sharma et al. [9] reported that rice ear head bug population was first recorded on 36<sup>th</sup> standard week as 1 nos./10 sweep nets and peak population (17 nos./10 sweep nets) was observed during 44<sup>th</sup> standard week.

### 3.4 Brown Plant Hopper (*Nilaparvata lugens*)

The infestation of brown plant hopper was recorded at weekly intervals. During *Kharif*, 2021, the initial population of (*N. lugens*) was recorded in 29<sup>th</sup> SMW (0.20 hopper per hill) and increase up to peak level of (11.25 hopper per hill) in 37<sup>th</sup> SMW. Thereafter, yellow brown plant hopper declined and it was not appeared 43<sup>th</sup> SMW. It was non-significant positive correlation with minimum temperature & RH (0.463 & 0.357) and negative correlation with maximum temperature & rainfall (-0.066 & -0.088) whereas, significant negative correlation with sunshine (-0.597\*), (Table 2 and Fig. 1). The present finding was closely associated with Patil et al. [10] reported that incidence of brown plant hopper, (*Nirparvata lugens*) (Stal.) was started from 33<sup>rd</sup> SMW i.e., second week of August with its peak during 41<sup>st</sup> SMW i.e., second week of October. These results are also in closed associated with Patil et al. [11] reported the incidence of brown plant hopper, *Nirparvata lugens* (Stal.) was started from 32<sup>nd</sup> SMW with its peak during 42<sup>nd</sup> SMW.

### 3.5 White Backed Plant Hopper (*Sogatella furcifera*)

The infestation of white backed plant hopper was recorded at weekly intervals during *Kharif*, 2021, the initial population of (*S. furcifera*) was recorded in 30<sup>th</sup> SMW (0.50 hopper per hill) and increase up to peak level of (12.80 hopper per hill) in 38<sup>th</sup> SMW. Thereafter, white backed plant hopper declined and it was not appeared 42<sup>th</sup> SMW. It was non-significant positive correlation with minimum temperature, RH & rainfall (0.300, 0.337 & 0.185) and negative correlation with maximum temperature (-0.300) whereas, significant negative correlation with sunshine (-0.094\*), (Table 2 and Fig. 1). These findings were

close association with to Sharma et al. [12] reported the first appearance of WBPH Started from 30<sup>th</sup> standard metrological week and reached to its peak during 40<sup>th</sup> standard week. These results are also in closed association with Verma et al. [13] incidence started from 33<sup>rd</sup> standard meteorological week and it attends its peak level during 38<sup>th</sup> standard meteorological week.

### 3.6 Green Leaf Hopper (*Nephotettix nigropictus*)

The infestation of green leaf hopper was recorded at weekly intervals during *Kharif*, 2021, the initial population of (*N. nigropictus*) was recorded in 41<sup>th</sup> SMW (0.20 hopper per hill) and increase up to peak level of (20.80 per cent per hill) in 37<sup>th</sup> SMW. Thereafter, green leaf hopper declined and it was not appeared 42<sup>th</sup> SMW. It was non-significant positive correlation with minimum temperature & RH (0.479 & 0.369) and negative correlation with maximum temperature & rainfall (-0.077 & -0.120) whereas, significant negative correlation with sunshine (-0.633\*), (Table 2 and Fig. 1). These findings were close association with to Sharma et al. [12] reported the first appearance of GLH Started from 30<sup>th</sup> standard metrological week and reached to its peak during 40<sup>th</sup> standard week. These results are also in closed association with Verma et al. [13] incidence started from 33<sup>rd</sup> standard meteorological week and it attends its peak level during 38<sup>th</sup> standard meteorological week [14-17].

## 4. CONCLUSION

The infestation of yellow stem borer (dead heart) was positive correlation with minimum temperature, maximum temperature, RH and negative correlation with rain fall, whereas significant negative correlation. The infestation of leaf folder was positive correlation with minimum temperature, RH and negative correlation with maximum temperature, rain fall and sunshine. The incidence of ear head bug was positive correlation with minimum temperature, RH, rainfall and sunshine whereas negative correlation with maximum temperature. The incidence of BPH was positive correlation with minimum temperature, RH, and negative correlation with maximum temperature & rainfall whereas significant negative correlation with sunshine. The incidence of WBPH was positive correlation with minimum temperature, RH, rainfall and negative correlation with maximum

temperature & sunshine. The incidence of GLH was positive correlation with minimum temperature, RH, and negative correlation with maximum temperature & rainfall whereas significant negative correlation with sunshine.

## ACKNOWLEDGEMENT

The authors are highly thankful to the Head, Department of Entomology, College of Agriculture, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (U.P.) India for providing essential facilities and support during the experimentation.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Shahbandeh M; 2019. Statista. Available:(<https://www.statista.com/statistics/271969/world-riceacreage-since-2008/>).
2. Anonymous. Directorate of Economics and Statistics, Department of Agriculture, Cooperation and Farmers Welfare; 2019.
3. Neeta J, Shukla BC, Gupta R, Agrawal RK. Seasonal abundance and activity of earhead bug in Chhattisgarh region of Madhya Pradesh. Ind. J. Ecol. 2013;14(1): 116-122.
4. Singh S, Kaur P, Kumar V, Singh H. Incidence of insect pest damage in rice crop in relation to meteorological parameters in Punjab—a plant clinic data based case study. Journal of Agrometeorology. 2012;14(1):50-3..
5. Jasrotia P, Khippal A, Yadav J, Kashyap PL, Kumar S, Singh GP. Effect of weather variables on the incidence of yellow stem borer (*Scirpophaga incertulas* W.) and leaf folder (*Cnaphalocrocis medinalis* G.) in rice. Journal of Cereal Research. 2019;11(3):247-251.
6. Patel S, Singh CP. Seasonal incidence of rice stem borer, *Scirpophaga incertulas* (Walker) on different varieties of rice in relation to weather parameters. J. Entomol. Zool. Stud. 2017;5(3):80-83.
7. Priya P, Prasad R. Population dynamics of rice leaf folder in relation to weather factors at Ranchi. Int. J. Manag. Sci. Eng. Manag. 2021;6(6):2456-1304.
8. Girish VP, Mahabaleshwar H, Hanamaratti NG, Balikai RA. Population dynamics of leafhopper, grasshopper, horned caterpillar and ear head bugs under different planting methods of rice ecosystem. Journal of Experimental Zoology, India. 2012;15(2): 451-454.
9. Sharma KR, Raju SVS, Singh KN, Roshan DR. Effects of environmental factors on population dynamics of rice earhead bug and their management with newer insecticide combinations and sole insecticide. Bangladesh Journal of Botany. 2019;48(4):973-979.
10. Patil SD, Patil HM, Pardeshi SR, Kusalkar DV, Sonawane KM, Bhoite KD. Seasonal incidence and impact of abiotic factors on pests of rice. The Pharma Innovation Journal. 2021;10(5):581-584.
11. Patil SD, Kusalkar DV, Patil HM, Bhoite KD. Seasonal incidence of insect pests on rice and impact of various abiotic factors on their incidence. Journal of Pharmacognosy and Phytochemistry. 2020;9(2):1869-1872.
12. Sharma KR, Raju SVS, Jaiswal DK. Influence of environmental effect on the population dynamics of brown plant hopper, *Nilaparvata lugens* (Stal) and white-backed plant hopper, *Sogatella furcifera* (Hovarth) in Varanasi region. Journal of Entomological Research. 2018;42(3):339-342.
13. Varma PK, Singh H, Singh B, Singh RP, Varma A. Population dynamics of brown plant hopper, *Nilaparvata lugens* (Stal.) in basmati rice and its correlation with abiotic factors in western plain zone of Uttar Pradesh. J. Pharm. Innov. 2021;SP-10(11):35-38.
14. Khare VK, Prakash R, Sneha, Kunal G. Effect of abiotic factors on the population dynamics of paddy earhead bug, *Leptocorisa oratorius* (F.). Journal of Entomology and Zoology Studies. 2020;8(6):157-160.
15. Kumar V, Sachan SK, Kumar S, Kumar A. Studies on the insect fauna and population dynamics of yellow stem borer and leaf folder in relation to abiotic factors in western Uttar Pradesh. J. Entomol. Zool. Stud. 2020;8(3):956-959.
16. Patel S, Singh CP. Seasonal incidence of rice stem borer, *Scirpophaga incertulas* (Walker) on different varieties of rice in



- relation to weather parameters. J. Entomol. Zool. Stud. 2017;5(3):80-83.
17. Sawai HR, Kothikar RB. Population dynamics of insect-pests of paddy and its correlation with weather parameters. International Journal of Chemical Studies. 2019;7(5):21-23.

---

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/100886>