



# Effect of Plant Density on Yield and Yield Components of Cowpea (*Vigna unguiculata* L.) in the Semi Arid Region of Brazil

Maria Eduarda Pereira dos Santos <sup>a++</sup>,  
Paulo César Ferreira Linhares <sup>a#\*</sup>,  
Thiffany Louhanne Alves Dantas <sup>a++</sup>,  
Lunara de Sousa Alves <sup>b†</sup>,  
Roberto Pequeno de Sousa <sup>c^</sup>, Uilma Laurentino da Silva <sup>c‡</sup>,  
Walter Martins Rodrigues <sup>c^</sup>, Joaquim Odilon Pereira <sup>c^</sup>,  
Pedro Ramualyson Fernandes Sampaio <sup>c##</sup>,  
Diego Zidane Fernandes da Costa <sup>a++</sup>,  
Débora Viviane Alves <sup>a++</sup>, Joyce Fernandes de Medeiros <sup>a++</sup>  
and Irene Leticia de Lima Cipriano Carlos <sup>a++</sup>

<sup>a</sup> Jitirana Research Group, Federal Rural University, Semi-Arid, Mossoró, RN, Brazil.

<sup>b</sup> Jitirana Research Group, Mossoró, RN, Brazil.

<sup>c</sup> Federal Rural University, Semi-arid, Mossoró, RN, Brazil.

## Authors' contributions

This work was carried out in collaboration among all authors. The execution of activities, such as planning, planting, conducting the experiment in the field, as well as writing, had the participation of all authors and was of great importance for the conclusion of the article. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajrcs/2024/v9i2274>

<sup>++</sup> Agronomy Engineering Student, Researcher and Member;

<sup>#</sup> PhD Researcher and Leader;

<sup>†</sup> PhD Researcher;

<sup>‡</sup> Agronomy Student;

<sup>^</sup> Professor;

<sup>##</sup> PhD Researcher of Phytotechnics;

\*Corresponding author: E-mail: paulolinhares@ufersa.edu.br;

**Cite as:** Santos, Maria Eduarda Pereira dos, Paulo César Ferreira Linhares, Thiffany Louhanne Alves Dantas, Lunara de Sousa Alves, Roberto Pequeno de Sousa, Uilma Laurentino da Silva, Walter Martins Rodrigues, Joaquim Odilon Pereira, Pedro Ramualyson Fernandes Sampaio, Diego Zidane Fernandes da Costa, Débora Viviane Alves, Joyce Fernandes de Medeiros, and Irene Leticia de Lima Cipriano Carlos. 2024. "Effect of Plant Density on Yield and Yield Components of Cowpea (*Vigna unguiculata* L.) in the Semi Arid Region of Brazil". *Asian Journal of Research in Crop Science* 9 (2):134-41. <https://doi.org/10.9734/ajrcs/2024/v9i2274>.

**Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116878>

**Original Research Article**

**Received: 27/03/2024**

**Accepted: 01/06/2024**

**Published: 07/06/2024**

## ABSTRACT

Cowpea cultivation is of great importance for farmers who work in this agricultural activity, in this sense, we evaluated the agronomic characteristics of cowpea (*Vigna unguiculata* (L.)) Creole cultivar (Semper verde), in the semi-arid region of Brazil.

The experiment was carried out from November 2021 to February 2022 at the Rafael Fernandes experimental farm, in the district of Alagoinha (5°03'37 "S, 37°23'50" W), northwest of Mossoró, State of Rio Grande do Norte, Brazil, with an area of 400 hectares.

A randomized complete block design was used with five treatments, which will consist of planting densities (35,714; 71,428; 107,142; 142,856 and 17,8570 plants ha<sup>-1</sup>) with four replications. Each plot consisted of dimensions of 5.0 m x 4.0 m, with six rows of 5.0 m in length, with a total area of 20.0 m<sup>2</sup>. The spacing used between rows was 0.70 m, while intra-row spacing was 0.40m.

The characteristics evaluated were: pod length, expressed in cm (PL), number of grains, expressed in pod<sup>-1</sup> units (NG), Pod weight, expressed in grams (PW), number of grains, expressed in pod<sup>-1</sup> units (NG), Weight of 100 grains, expressed in grams (P100G) and grain production, expressed in kg ha<sup>-1</sup> (PG) of cowpea, Creole cultivar (Sempre verde). Planting densities contributed greatly to cowpea productivity, with productivity of 184.82 kg ha<sup>-1</sup>, at a density of 142,856 plants ha<sup>-1</sup>.

The cowpea cultivar (Semper Verde) is agronomically viable for planting in the semi-arid region.

**Keywords:** Family farming; agroecological production and grain production; planting; cowpea.

## 1. INTRODUCTION

Cowpea [*Vigna unguiculata* (L) Walp.] is a legume of great importance for family farmers, considering that it contributes to the social, nutritional and economic aspects of thousands of people [1]. In the northeast region of Brazil, this crop is planted mainly during the rainy season, when farmers cultivate large areas to produce grains for food and commercialization, bringing income to those who work in this activity.

It is worth noting that this crop has genetic variability, and can be used for different purposes, being sold in different ways, mainly as dry grain, green grain, green pod and seeds, greatly contributing to increasing the producer's income [2]. The low technological level, associated with irregular rainfall, predisposes bean cultivation to low productivity in the northeast region of Brazil, making it especially important for producers to cultivate crops during the rainy season in the region.

In this sense, it is extremely important to use technologies that will contribute to an increase in cowpea productivity in the northeast region of Brazil, which will bring economic returns to farmers in the region [3].

To achieve the maximum productive potential of cowpea, it is extremely important to consider the number of plants per unit area, influenced by the morphological, physiological and grain yield characteristics.

The use of cowpea in the northeast region of Brazil constitutes an extremely important alternative for those who work in this activity, requiring the development of the crop in edaphic-climatic conditions to understand the behavior of the crop and its maximum agronomic efficiency.

It is worth noting that this culture is extremely important for the semi-arid region, however, the lack of knowledge regarding technology transfer is present in many areas [4].

In view of the above, we evaluated the agronomic characteristics of cowpea (*Vigna unguiculata* (L.)) Creole cultivar (Semper verde), in the semi-arid region of Brazil.

## 2. MATERIALS AND METHODS

### 2.1 Location of the Experiment Installation

The experiment was conducted from September to December 2021 at the Rafael Fernandes experimental farm, in the district of Alagoinha (5°03'37" S, 37°23'50" W), northwest of Mossoró, State of Rio Grande do Norte, Brazil, in the period November 2021 to February 2022. According to [5] and the Köppen classification, the local climate is BSw<sup>h</sup>, dry and very hot, with a dry season, often from June to January, and a rainy season from February to May, average annual precipitation of 673.9 mm and average relative humidity of 68.9%.

The region's climate, according to the Köppen Geiger classification, is dry and very hot with two seasons: a dry season starting in June and ending in January and a rainy season starting in February and ending in May [5]. The soil in the experimental area was classified as a typical dystrophic red yellow argisol, with a sandy loam texture [6].

Before setting up the experiment, soil samples were collected from the 0-20 cm depth, homogenized, and sent to the soil chemistry and fertility laboratory for the following analyses: soil pH = 6.6; EC = 0.44 dS m<sup>-1</sup>; O.M. = 9,8 g kg<sup>-1</sup>; N = 0.52 g kg<sup>-1</sup>; P = 44.30 mg dm<sup>-3</sup>; K = 60.28 mg dm<sup>-3</sup>; Ca=18.50 cmolc dm<sup>-3</sup>; Mg = 6.80 cmolc dm<sup>-3</sup>; Na = 2.33 mg dm<sup>-3</sup>; Cu = 0.60 mg dm<sup>-3</sup>; Fe = 4.30 mg dm<sup>-3</sup>; Mn = 10.45 mg dm<sup>-3</sup>; and Zn = 4.56 mg dm<sup>-3</sup>.

### 2.2 Experimental Design

A randomized complete block design was used with five treatments, which will consist of planting densities (35,714; 71,428; 107,142; 142,856 and 17,8570 plants ha<sup>-1</sup>) with four replications, and the cultivar used was Creole Semper Verde. Each plot consisted of dimensions of 5.0 m x 4.0 m, with six rows of 5.0 m in length, with a total area of 20.0 m<sup>2</sup>. The spacing used between rows was 0.70 m, while intra-row spacing was 0.40m.

During the period in which the cowpea crop was in the field, weeding was carried out to remove invasive plants that compete for water and nutrients during the crop cycle. Irrigation was

carried out by drip, with a daily irrigation shift divided into two applications (morning and afternoon).

When harvesting took place, the cowpea pods were taken to the Phytotechnics Laboratory of the Department of Agricultural and Forestry Sciences belonging to UFERS to evaluate the following characteristics: pod length, expressed in cm (PL), number of grains, expressed in pod<sup>-1</sup> units (NG), Pod weight, expressed in grams (PW), number of grains, expressed in pod<sup>-1</sup> units (NG), Weight of 100 grains, expressed in grams (P100G) and grain yield, expressed in kg ha<sup>-1</sup> (PG) of cowpea, Creole cultivar (Sempre verde).

### 2.3 Statistical Analysis

Statistical analysis was performed according to conventional methods of analysis of variance [7], using ESTAT statistical software [8]. The response curve fitting procedure was performed using the ESTAT Software [8], applying regression analysis and conducting hypothesis testing that helps the researcher accept or reject a statistical hypothesis based on experimental results [9,10].

## 3. RESULTS AND DISCUSSION

It was observed that there was a significant? Difference for all cowpea characteristics evaluated at the level of p< 0.01 probability (Table 1). For the characteristics of pod length, numbers of pod grains, pod weight and weight of 100 grains, the data were adjusted to the linear equation (Figs. 1 to 4), in relation to grain production the data were adjusted to a quadratic equation (Fig. 5). The use of landraces of cowpea is extremely important as an alternative for farmers working in family farming, reducing production costs and increasing profitability [11].


These variables are extremely important, as they demonstrate the productive aspect of cowpea, being used by farmers as an agronomic parameter.

Pod length showed the greatest growth in density of 35,714 plants ha<sup>-1</sup>, with a value of 19.46 cm, 3.85 cm higher than the highest planting density (178,570 plants ha<sup>-1</sup>) (Fig. 1). Pod length is an extremely important characteristic, as it is related to the number of pod<sup>-1</sup> seeds. [12], evaluating cowpea productivity under doses and sources of organic fertilizer in the municipality of Chapadinha-MA, found a pod length of 20.5 cm with the application of 20 t ha<sup>-1</sup>, higher than the aforementioned research.


**Table 1. F values for pod length, expressed in cm (PL), number of grains, expressed in pod<sup>-1</sup> units (NG), Pod weight, expressed in grams (PW), number of grains, expressed in pod<sup>-1</sup> units (NG), Weight of 100 grains, expressed in grams (P100G) and grain production, expressed in kg ha<sup>-1</sup> (PG) of cowpea, Creole cultivar (Sempre verde)**

Causes of Variation	GL	PL	NG	PW	NG	P100G	PG
Blocks	3	11.08**	9,30**	7.65**	10.78**	8,45**	6,49**
Treatments	4	16.47**	8.16**	17.02**	14.18**	10.27**	7.46**
Residue	12	---	---	---	---	-----	-----
CV (%)	---	13.42	10.12	14.75	23.1	6.17	5.49

\*\* = P<0,01; \* = P<0,05; ns = not significant



**Fig. 1. Pod length as a function of different planting densities in the semi-arid region of Brazil**



**Fig. 2. Number of grains as a function of different planting densities in the semi-arid region of Brazil**

According to [13], the number of grains per pod is directly related to the length of the pods, thus explaining the results found in this research, as there was little variation in the size of the pods and also no statistical difference.

The highest number of grains per pod was observed at a density of 35,714 plants ha<sup>-1</sup>, with

a maximum value of 14.65 pod units<sup>-1</sup> (Fig. 2). The number of pod<sup>-1</sup> grains is of great importance, as it is related to productivity. [14], evaluating cowpea production depending on the use of inoculant and organic and mineral fertilizers, found a number of pod<sup>-1</sup> grains of 14.3 units with biofertilizer application, which is lower than the aforementioned research. [15]

evaluating yield of Pingo de Ouro cowpea bean under different sowing densities and goat manure doses in the region of Mossoró, Brazil, with a number of seeds worth 12.0 pod units<sup>-1</sup>, was lower than the density of 35,714 plants ha<sup>-1</sup>.

For pod weight, there was a decrease depending on planting densities, with a maximum value of 5.9 g at a density of 35,714 plants ha<sup>-1</sup> (Fig. 3). Pod weight is an important characteristic, as it is related to the productive aspect of the crop. [16], evaluating the characteristics and pod yield of snap beans depending on sources and doses of organic matter, found a pod weight of 12 g at a dose of 10 t/ha, a value higher than that of the aforementioned research. [17], studying the characterization of yield in green cowpea grains, found a pod weight of 11.08 g, higher than the aforementioned research.

In relation to the number of grains per pod, the density of 35,714 plants ha<sup>-1</sup> was the one that promoted the greatest increase in the characteristic, with a maximum value of 14.7 units per pod. In subsequent densities, there was a decrease to a density of 178,570 plants with a value of 11.9 grain units per pod (Fig. 4). [18], studying the productivity and water use efficiency of cowpea cultivars under drip irrigation, found a number of grains per pod of 11.1 units, a value lower than that of the aforementioned research. [19], studying the potential of cowpea for the green pods and grains market, found an average number of green grains under cultivation conditions (rainfed) of 15.1 units per pod, which is lower than the aforementioned research, a value close to that of the aforementioned research.




Fig. 3. Pod weight as a function of different planting densities in the semi-arid region of Brazil






Fig. 4. Number of grains as a function of different planting densities in the semi-arid region of Brazil



**Fig. 5. Weight of 100 grains as a function of different planting densities in the semi-arid region of Brazil**



**Fig. 6. Grain production as a function of different planting densities in the semi-arid region of Brazil**

The mass of 100 grains is a very important characteristic, as it is related to the productive variable, and in this research, the density of 35,714 plants ha<sup>-1</sup> obtained the highest average mass, with a value of 24.81 g (Fig. 5). In higher densities there was a decrease, with a minimum weight in density of 178,570 plants ha<sup>-1</sup>, with a value of 21.3 g. [20], studying the productivity and water use efficiency of cowpea cultivars under drip irrigation, found a weight of 100 grains of 20.22 g, a value lower than that of the aforementioned research.

The highest grain production was obtained at a density of 142,856 plants ha<sup>-1</sup>, with a maximum value of 184.82 kg ha<sup>-1</sup>, with an increase depending on the other densities, with a value of 164.10 kg ha<sup>-1</sup> (Fig. 6). The observed productivity is a function of the Creole cultivar

used. [21], studying the productivity and water use efficiency of cowpea cultivars under drip irrigation, found a weight of 100 grains of 20.22 g, a value lower than that of the aforementioned research. [22], studying the morphological and agronomic characterization of cowpea accessions collected in municipalities in Rio Grande Norte, Brazil, found an average productivity of dry pods in the second harvest of accessions A323, AC10 and AC22 of 286.51, 324.41 and 355.90 kg ha<sup>-1</sup>, values higher than those in the aforementioned research. [23] studying the production of semi-prostrate cowpea in rainfed and irrigated crops with an average productivity of 851.59 kg ha<sup>-1</sup>, higher than the aforementioned research.

It is worth mentioning that the size of the grain, as well as the color, constitutes a market

preference and is important in determining the price of the product, being characters that should not be markedly altered during the selection process, since there is a preference for grains with a weight of 100 grains around 18 g of kidney-shaped or rounded shapes [24]. Knowledge of the association of grain productivity and its components is important for the selection of parents and in promising segregating populations, making the selection process efficient [25].

#### 4. CONCLUSION

Planting densities contributed greatly to cowpea productivity, with productivity of 184.82 kg ha<sup>-1</sup>, at a density of 142,856 plants ha<sup>-1</sup>. The cowpea cultivar (Semper Verde) is agronomically viable for planting in the semi-arid region. The low technological level, associated with irregular rainfall, predisposes bean cultivation to low productivity in the northeast region of Brazil, making it especially important for producers to cultivate crops during the rainy season.

#### ACKNOWLEDGEMENTS

To the Jitirana research group (CNPQ-National Council for Scientific and Technological Development, Brazil), a pioneer in Brazil in the use of spontaneous species from the semi-arid region as green manure in the agroecological production of leafy and root vegetables, and the Universidade Federal Rural do Semi-Árido-UFERSA, for the physical structure and granting of scientific initiation scholarships for the development of research.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Oliveira IJde, Fontes JRA, Dias MC, Barreto JF. Technical recommendations for the cultivation of cowpea in the State of Amazonas. (Technical Circular, 71). Manaus: Embrapa Amazônia Oeste; 2019.
2. Ramos HMM, Bastos EA, Andrade Júnior AS, Cardoso MJ. Water use efficiency and cowpea grain productivity under different water regimes. In: National Cowbean Congress, 3. Recife-PE. Annals... CONAC, Recife/PE; 2013.
3. Alves NFdeO, Linhares PCF, Sousa VEde, Carlos KGdaS, Santos MdeFAdos, Peixôto, LSdeL, Lobato LVC, Leite IdeO, Paiva MRdaS, Assis JPde, Rodrigues WM, Lima IRPde. Productivity of Dry Grains of Cowpea [*Vigna unguiculata* (L.) Creole Cultivar (Canupum), as a Function of Different Planting Densities. Asian Journal of Research in Crop Science. 2023;8(4): 421–428. Available:<https://doi.org/10.9734/ajrcs/2023/v8i4222>
4. Vale JCdo, Bertini CBA. Cowpea: From planting to harvest. Viçosa. 2017;3(1). Available:<https://doi.org/10.33447/paubrasilia.v3i1>
5. Beck HE, Zimmermann NE, Mcvigar TR, Vergopolan N, Berg A, Wood EF. Present and future Köppen-Geiger climate classification maps at 1-km resolution. Scientific Data. 2018;5:180214.
6. Santos JAS, Soares CMG, Corrêa AM, Teodoro PE, Ribeiro LPE, Abreu HKA. Agronomic performance and genetic dissimilarity among cowpea (*Vigna unguiculata* (L.) Walp.) genotypes. Global Advanced Research Journal of Agricultural Science. 2014;3(1):271-277.
7. Banzatto DA, Kronka SN. Agricultural experimentation. 3rd ed. Jaboticabal: FUNEP. 1995:245.
8. Barbosa JC, Malheiros EB, Banzatto D. A. ESTAT: A system for statistical analysis of agronomic trials. Jaboticabal: Unesp, Version 2.0; 1992.
9. Assis JP, Sousa RP, Linhares PCF. Statistical hypothesis testing. EdUFERSA; 2020. Available:<https://livraria.ufersa.edu.br/wp-content/uploads/sites/165/2020/08/testes-de-hipoteses-estatisticas-edufersa.pdf>
10. Assis JP. Simple linear regression, simple linear correlation, multiple linear regression and multiple linear correlation. EdUFERSA. 2013;310. Available:<https://livraria.ufersa.edu.br/regressao-e-correlacao-linear-simples-e-multiplica/>
11. Abreu L, Cansi E, Juriatti C. Assessment of the socioeconomic performance of landraces and commercial corn hybrids in the Chapecó microregion. Brazilian Journal of Agroecology. 2007;2(1):1230-1233.
12. Oliveira CAB, Mello Pelá G, Pelá A. Inoculation with *Rhizobium tropici* and foliar fertilization with molybdenum in

- common bean crops. Journal of Neotropical Agriculture. 2017;4(5):43-50.
13. Cavalcante EdaS, Freire Filho FR, Rocha MdeM, Goes ACP, Ribeiro VQ, Silva KJD. BRS Tumucumaque: Cowpea Cultivar for Amapá and Other States of Brazil. Macapá: EMBRAPA Amapá. 2014;5. (Technical Announcement 124).
  14. Silva CF, Moura MFde, Vilela Ávilo RR, Araújo MBde, Marques JDS. Production of cowpea depending on the use of inoculant and organic and mineral fertilizers. Diversitas Journal. 2019;4(3):1130–1145. Available:<https://doi.org/10.17648/diversitas-journal-v4i3.832>
  15. Silva RASda, Linhares PCF, Souza ASdos, Pereira MFS, Assis JPde, Sousa RPde, Neves AM, Alves LSde, Silva RIGda. Yield of pingo de ouro cowpea bean under different sowing densities and goat manure doses in the region of Mossoró, Brazil. Journal of Agricultural Science. 2018;10(12):329-335. Available:<https://ccsenet.org/journal/index.php/jas/article/view/0/37346>
  16. Santos GM, Oliveira AP, Silva JAL, Alves EU, Costa CC. Characteristics and pod yield of snap beans depending on sources and doses of organic matter. Brazilian Horticulture, Brasília. 2001;19(1):30 - 35.
  17. Coelho TJSda, Coelho RRP, Santos VS. Characterization of yield in green cowpea grains. IV National Agroindustry Meeting; 2018.
  18. Pimenta LJL, dos Santos SR, Bernardino DLMP, Barbosa JAE, Alves AGT, De Carvalho AJ. Productivity and water use efficiency of cowpea cultivars under drip irrigation. Contributions to Social Sciences. 2023;16(11):26504–26522. Available:<https://doi.org/10.55905/revconv.16n.11-104>
  19. Carvalho CDM, Francelino FMA, Manhães CMC, Silva MPS, Ribeiro EC, Bueno JAR. Rendimento de grãos secos e componentes de produção de cultivares de feijão-caupi (*Vigna unguiculata* L.) Em cultivo irrigado na estação da seca em Pedro Afonso – To. Jornada de Iniciação Científica e Extensão. Instituto Federal do Tocantins. 2020;1-8.
  20. Pimenta LJL, dos Santos SR, Bernardino DLMP, Barbosa JAE, Alves AGT, De Carvalho AJ. Productivity and water use efficiency of cowpea cultivars under drip irrigation. Contributions to Social Sciences. 2023;16(11):26504–26522. Available:<https://doi.org/10.55905/revconv.16n.11-104>
  21. Pimenta LJL, dos Santos SR, Bernardino DLMP, Barbosa JAE, Alves AGT, De Carvalho AJ. Productivity and water use efficiency of cowpea cultivars under drip irrigation. Contributions to Social Sciences. 2023;16(11):26504–26522. Available:<https://doi.org/10.55905/revconv.16n.11-104>
  22. Freitas SQ. (Agronomic characterization of cowpea accessions collected in municipalities in the state of Rio Grande do Norte. 2010. 39f. Monograph (graduation) – Universidade Federal Rural do Semi-Árido (UFERSA), Mossoró; 2010.
  23. Silva Alda, Neves JA. Production of semi-prostrate cowpea in rainfed and irrigated crops. Brazilian Journal of Agricultural Sciences, 6(1): 29-36.
  24. Saraiva KR, Oliveira KR, Marques Filho F, Silva Fs, Silva FS, Sales JRS. Productive aspects and biomass of cowpea (*Vigna unguiculata*) under doses of bovine biofertilizer in agroecological cultivation. Family Agriculture: Research, Training and Development. 2020;14(1): 184-198.
  25. Benvindo RN, Silva JALda, Freire Filho FR, Almeida ALGde, Oliveira JTS, Bezerra AAC. Evaluation of semi-prostrate cowpea genotypes in rainfed and irrigated cultivation. Comunicata Scientiae. 2010; 1(1):23-28.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/116878>