



Variation of the Intensity of the Gamma Radiation in the Period from May 28 to September 25 of 2017 in Sao José dos Campos, SP, Brazil

Inacio M. Martin^{1*}, Franklin A. Silva¹ and Marcelo P. Gomes¹

¹*Technological Institute of Aeronautics (ITA), SP, Brazil.*

Authors' contributions

This work was carried out in collaboration between all authors. Author IMM designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors MPG and FAS managed the analyses of the study. Author FAS managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/PSIJ/2017/37789

Editor(s):

- (1) Manuel Malaver de la Fuente, Universidad Maritima del Caribe, Av. El Ejercito, Catia La Mar, Estado Vargas, Venezuela.
(2) Roberto Oscar Aquilano, School of Exact Science, National University of Rosario (UNR), Rosario, Physics Institute (IFIR)(CONICET-UNR), Argentina.

Reviewers:

- (1) Samuel Sonko Sackey, University of Cape Coast, Ghana.
(2) José Martínez Reyes, University of the Ciénega of Michoacán State, México.
(3) Krzysztof Kozak, Institute of Nuclear Physics Polish Academy of Sciences, Poland.
(4) Adriana Estokova, Technical University of Kosice, Slovakia.
Complete Peer review History: <http://www.sciencedomain.org/review-history/22516>

Original Research Article

Received 28th October 2017
Accepted 22nd December 2017
Published 29th December 2017

ABSTRACT

The variation of gamma radiation intensity between 200 keV to 10.0 MeV was measured, in the period from June 28 to September 25 of 2017. These measurements were taken 25 meters above ground level in a tower in São José dos Campos, SP, Brazil. A minute interval was used from each point of measurements. During this period, there were weak and moderate rains with net intensity of 27 mm. The variation in temperature during this period was between 15°C and 32°C, reproducing a desert-like climate. By monitoring the gamma radiation it was possible to observe the arrival of cold fronts from Southern Brazil and the (day/night) cycles due to the greater or lesser amount of radon gas present in the region. The main feature of this work is that it indicates the dynamics of gamma radiation in a simple way according to the variation of meteorological parameters in that location, which is very important for environmental studies in the tropical region.

*Corresponding author: E-mail: martin@ita.br;

Keywords: Gamma radiation; cold fronts; radon gas; meteorological parameters; weak and moderate rains; day/night cycles.

1. INTRODUCTION

At the (ground / air) interface of Earth's surface, the ionizing radiation is composed mainly of radon gas, the telluric radiation of the soil and the radiation of the primary and secondary cosmic rays. However, it is difficult to separate over time the intensity of the ionizing radiation emanating from each component as the energies overlap. The telluric radiation is given by ^{238}U , ^{235}U , ^{40}K , ^{232}Th and is constant for each region [1]. The radon gas that comes from the disintegration of ^{238}U of Earth's crust [2] into Ra-226 to Rn-222 arriving at the isotopes ^{214}Pb , ^{214}Po and ^{214}Bi giving alpha and gamma radiation. The primary cosmic radiation consists mainly of galactic and extragalactic protons and from the Sun with very high energy that interacts with Earth's atmosphere producing the EAS (Extensive Air Showers) [3]. The efficiency of this interaction is maximum when it occurs at altitudes between 15 and 17 km in the tropics, which form secondary cosmic rays with muonic, mesonic, and neutronic components that reach the Earth's surface in the region [4]. These radiations can cause health problems for the crew and passengers of civil/military aviation in this altitude that are present at the beginning of the stratosphere called Pfozter maximum. However, this component contributes less to the concentration of ionizing radiation on the Earth's surface. Another possible source of ionizing radiation in the Earth's lower atmosphere is produced by electrical discharges between cloud-earth, earth-

cloud and cloud-cloud. X-rays, gamma rays, neutrons and beta particles are all formed by the lightning cone [5]. Other sources of ionizing radiation are those produced in industry, medical, dental clinics and hospitals, but these radiations are mainly controlled in small and specific areas.

2. MATERIALS AND METHODS

The gamma ray detector for the energy range of 200 keV to 10.0 MeV consists of a 3-inch-by-3-inch-diameter and high sodium iodide scintillation crystal (3" x 3"), doped with thallium. This crystal is directly coupled to a photomultiplier (PM), which registers the pulses coming from the scintillator and with amplification and an analog digital converter (ADC) these digitized signals are recorded by a computer [6]. The detector was in a closed room and air exchange was ensured during the measurements. The experimental set (Fig. 1) is located in the inner room of a tower (23°12'45.0"S 45°52'00.3"W), 25 meters high in relation to the ground surface.

The scintillator coupled to photomultiplier is wrapped in a thin layer of aluminum to make it portable. Scintillator and associated electronics were calibrated in terms of energy and counting intensity per minute, at the laboratory of experimental teaching physics of ITA (Technological Institute of Aeronautics), using radioactive sources and a spectral analyzer of counts versus energy in the range of 0.2 to 10 MeV (Million electron Volt), [7,8].



Fig. 1. View of the gamma scintillator associated with electronics and computer

Source: Project Atmosrad 2017 [9]

3. RESULTS AND DISCUSSION

Gamma radiation measurements were carried out during the period of June 28 to September 25 of 2017, in the inner room above the tower, seen in Fig. 2. During the interval described above, on the roof of the tower was placed the rain gauge that reported the intensity of rains in (mm/min).



Fig. 2. Exterior view of the tower with the room 25 meters above the ground level

Source: Project Atmosrad 2017 [9]

Fig. 3 shows the series of measured gamma radiation intensity between June 28 to September 25 of 2017, with uninterrupted monitoring minute by minute during this net time.

The smooth tool was used to transform data from minutes to days as seen on the green line. Analyzing the dynamics of the radiation measurements, there are 4 large variations occurring in the whole period analyzed. Between




Fig. 3. Monitoring of gamma radiation series between the start time and 130 x 10³ minutes after

Source: Project Atmosrad 2017 [9]

the beginning of monitoring and close to 70 x 10³ minutes, the mean intensity of the measured radiation was 37.5 x 10³ (counts / minute). In this period it was possible to see small variations which indicate passages of cold fronts without rain. See this dynamic in the expanded period in Fig. 4, taken from the graph in Fig. 3.

Fig. 5 shows the radiation monitoring between 70 to 80 x 10³ minutes after the start of the measures. It was a rainy week with intensities varying according to Fig. 5 and Fig. 8.

In the beginning between 70 and 71 x 10³ minutes there was intense rain, where the level of radiation count reached the order of 40 x 10³ (counts/min). On the other days, there were always less intense rains as shown by the gamma radiation peaks caused by the rains. In Fig. 6, taken during the measurement time of 80 to 100 x 10³ minutes, there are variations in the dynamics of the radiation with passages of two cold fronts in the region, but without causing rains. However, the terrestrial surface was wet and with very little exhalation of radon gas. The arrival of the cold front causes an increase in the radiation due to the accumulation of radon gas that arrives with these fronts.

In Fig. 7, the gamma radiation intensity measured between the times of 100 to 130 x 10³ minutes with average intensity of 37.3 x 10³ (counts / min), undergoes an influence of soil in the region that was very dry and high pressure that was 970 mbar when local background value is 940 mbar. This occurs in the afternoon where the temperature varied between 20° to 30°C and the night between 12° to 20°C. This dynamics in

temperature during this period facilitates greater and lesser exhalation of the radon gas, as shown in the Fig. 7 with cycles of exactly 24 hours. The

two largest radiation peaks, shown in Fig. 7, are caused by heavy fog in the mornings of those days.




Fig. 4. Monitoring of gamma radiation between the start time and 70×10^3 minutes after
 Source: Project Atmosrad 2017 [9]




Fig. 5. Monitoring of gamma radiation between 70×10^3 and 80×10^3 minutes after
 Source: Project Atmosrad 2017 [9]




Fig. 6. Monitoring of gamma radiation between 80×10^3 and 100×10^3 minutes after
 Source: Project Atmosrad 2017 [9]




Fig. 7. Monitoring of gamma radiation between 100×10^3 and 130×10^3 minutes after
 Source: Project Atmosrad 2017 [9]




Fig. 8. Spectrum in time of the rain that occurred in the region during this period of measurements
 Source: Project Atmosrad 2017 [9]

Fig. 8 shows the rainfall spectrum in (mm / min), varying in time. During the whole period, only 27 mm of net rain accumulated in the region in the course of a week.

“Historically, the mean rainfall value for the period mentioned above is around 170 mm”.

4. CONCLUSION

In the period of June to September of 2017, the intensity of rain was monitored every minute and in the same place. In the same time the intensity of low energy gamma rays was also measured. The analysis show that during one single week of

moderate and light rains, there was a noticeable increase in the intensity of gamma rays. The net rainfall in the period measured was 27 mm scattered in time. Analysis of all figures plotted in this work shows clearly the difference caused by the rains in the measurement of gamma rays. Also in this work, the perfect oscillation of the gamma rays (day/night) in the dry period is evidenced, without clouds, fog or lightning. This oscillation is caused by the exhalation of radon gas (Rn-222) in the region that is larger during the local solar zenith time with dry soil. The periodicity is partially or totally destroyed with the presence of fog, continuous rain and drizzle in the region.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gomes Marcelo Pego, Martin Inácio Malmonge, Silva Franklin Andrade, Sismanoglu Bogus Nubar. Monitoring of gamma radiations and meteorological parameters at ground level in São José dos Campos, Brazil. *Impact: International Journal of Research in Engineering and Technology*. 2016;4:8.
2. Martin IM, Douglas Carlos Vilela, Marcelo P. Gomes. Dynamics in times of ionizing radiation and rainfalls in tropical region of Brazil. *Asian Review of Environmental and Earth Sciences*. 2017;4:7-11.
3. Gusev AA, Jayanthi UB, Martin IM, Pugacheva GI, Spjeldvik WN. Nuclear reactions on rarest atmosphere as a source of magnetospheric positron radiation belt. *J. Geophys. Res.* 2001; 106(A11):26111–26116.
4. Gusev Anatoli A, Martin Inacio M, Alves Mauro A, de Abreu Alessandro. J. simulation of the radiation fallout from gamma-ray measurements. *Modeling Earth Systems and Environment*. 2015;1: 18.
5. Martin Inácio Malmonge, Gomes Marcelo P. Intensity variation of gamma radiation on ground level interface in São José dos Campos, SP, Brazil. *Environmental Science: An Indian Journal*. 2013;879-82.
6. Silva FA, Martin IM, Gomes MP. Monitoring of ionizing radiation and rain intensity during May to October 2015 in São José dos Campos, Brazil. *Davi: Journal of Environmental Science and Engineering*; 2016.
7. Gomes MP, Martin IM. Simultaneous measurements of rainfall intensity, low energy neutrons and gamma radiation in São Jose dos Campos, SP, Brazil. *Journal of Environmental Science and Engineering*. 2013;9:161-167.
8. Martin IM, Marcelo P. Gomes, Bogos Nubar Sismanoglu, Nicolas Cruvinel Lindo. Daily variability of radon gas in Brazilian tropics near ground level surface. *Journal of Environmental Science and Engineering*. 2015;A4:516-521.
9. Project Atmosrad. Research Group of CNPq. Available:<http://dgp.cnpq.br/dgp/espelhogrupo/4663704786429697>

© 2017 Martin et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/22516>