



Evaluation of Antibacterial Activity of Methanolic Extract of *Avicennia marina* Leaf Extract against *Staphylococcus aureus*

Swetha Ramasubramanian ^a≡, Anitha Roy ^{b*}⊙, Pitchiah Sivaperumal ^b⊙ and T. Lakshmi ^b≡

^a Saveetha Dental College and Hospitals, Saveetha institute of Medical sciences and Technical Sciences (SIMATS), Chennai - 600077, Tamil Nadu, India.

^b Department of Pharmacology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai - 600077, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i61A35587

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77720>

Original Research Article

**Received 20 October 2021
Accepted 26 December 2021
Published 28 December 2021**

ABSTRACT

Introduction: Mangrove are widespread in the tropical and subtropical regions of Indo-West-Pacific area. *Avicennia marina* possesses vast medicinal values. The pharmacological activity of the plant is attributed to the presence of several phytochemicals.

Aim: To evaluate the antibacterial activity of *Avicennia marina* leaf extract against *Staphylococcus aureus*.

Methods and Materials: Plant extract prepared by adding 20g of dried powdered mangrove leaf sample to 100 ml of methanol. MDRSA, MRSA & VRSA were cultured in Muller –Hinton Broth for 24 hr at room temperature. From this prepared bacterial suspension, 1ml of was spread over Muller Hinton agar plate and incubated for 24hrs at ambient temperature. Antibacterial activity carried out through the disc diffusion method. Whatman filter paper discs (5mm) were impregnated with various concentrations (50, 100, 150, 200, 250 & 300 µg/ml) of leaf extract. After incubating the plates for 24hr at room temperature the zone of inhibition was measured. Minimum Inhibitory

≡ Professor;

⊙ Assistant Professor;

*Corresponding author: E-mail: anitharoy@saveetha.com;

Concentration was determined in 5 concentrations (50 -300 µg/ml) with blank (extract in Muller Hinton broth). The inoculated bacteria in test tubes are incubated for 24hr in ambient temperature. The results are noted as well growth (+) and inhibited (-). Tetracycline (1µg/ml) was used as standard and DMSO as negative control.

Results: The zone of inhibition was greater at 300µg/ml of the extract for MDRSA (13±1.2) , MRSA ((13±1.4))& VRSA (9±1.2). MIC for MDRSA, MRSA and VRSA was 250µg/ml, 200µg/ml 150µg/ml respectively.

Conclusion: The present study showed effective antibacterial activity of *Avicennia marina* against MDRSA, MRSA, VRSA. Hence, this extract may be used for infections against resistant *Staphylococcus aureus*.

Keywords: *Avicennia marina*; antibacterial activity; mangroves; natural source; *Staphylococcus aureus*.

1. INTRODUCTION

Plants are recently explored for their pharmacological activities in different health conditions such as cancer, diabetes, inflammatory as well as oxidative stress conditions [1–10]. Mangroves are a group of trees and shrubs that live in the coastal intertidal zone. Mangrove trees can grow under adverse environmental conditions like low-oxygen soil, high temperature, high salinity, etc. The distribution of mangrove forests is mainly at the tropical and subtropical latitudes [11].

Mangrove plants are a rich source of steroids, triterpenes, saponins, flavonoids, alkaloids, tannins [12,13]. Extracts from different mangrove plants are reported to possess diverse medicinal properties such as antibacterial, anthelmintic [14]. Marine organisms including coastal plants and mangrove trees are known for their ability to produce secondary metabolites that differ from the terrestrial plants due to their specific adaptation such as aerial roots, viviparous seedlings, high tolerance to extreme temperature, weather and salinity [15-16]. Mangrove ecosystems serve as a natural filter of land-derived waste materials thus supporting the health of marine ecosystems [17- 19].

The *Avicennia marina* (Forssk.) Vierh., a mangrove tree belonging to the Acanthaceae family, is mostly found in the subtropical and tropical regions of the Indo-West-Pacific area. It is considered a representative example of mangroves that have been widely investigated for their medicinal importance [20]. The presence of the various categories of phytochemicals makes this plant an excellent candidate for the treatment of various health ailments. *Avicennia marina* has been used in the traditional medicine system for many centuries [21-24]. For example,

leaves are used for the treatment of ulcers, abscesses, and rheumatism. The leaf decoction is also used for the treatment of malarial fever and food poisoning [25]. Our team has extensive knowledge and research experience that has translated into high quality publications [26–31]. The aim of this study was to evaluate the antibacterial activity of *Avicennia marina* leaf extract against *Staphylococcus aureus*.

2. METHODS AND MATERIALS

2.1 Study Setting

Marine Biomedical and Environmental Health Research Lab - Blue Lab, Saveetha Dental College. The study was ethically approved by the Scientific Review Board, Saveetha Dental College, Chennai.

2.2 Collection of Plant Material and Preparation

The fresh leaves of *Avicennia marina* were collected from Tuticorin, Tamil Nadu. The leaves were washed thoroughly with tap water then shade dried on table tissue paper for 2-3weeks and turned into a fine powder.

2.3 Preparation of Extraction

20g of dried powdered mangrove leaf samples were mixed with 100ml of methanol/Ethanol and allowed to place for 24 hours at ambient temperature. Then the mixture was passing through whatman filter paper (No.4) then the filtrate was centrifuged at 3000rpm for 10min and further filtered by 0.45µm syringe micro filter. At last, the solvents are evaporated via vacuum rotary evaporator until samples are obtained in powder form. Then the sample was stored in a shadowy aluminum container at 4°C for further analysis.



Fig. 1. Collection of *Avicennia marina* plant collection and preparation

2.4 Bacterial Suspension

Multi drug resistant *Staphylococcus aureus* (MDRSA), Methicillin resistant *Staphylococcus aureus* (MRSA) and Vancomycin resistant *Staphylococcus aureus*(VRSA) was collected from Department of Microbiology, Saveetha Medical College and Hospital, Tamil Nadu. The bacterial pathogens were cultured in Muller–Hinton Broth for 24 hr at room temperature. From this bacterial suspension was prepared with saline and the optical density was measured at 600 nm. The concentration of microbial suspension was fixed as 10^8 CFU/ml. 1ml of suspension was spread over on Muller Hinton agar plate and incubated for 24hrs at ambient temperature.

2.5 Antibacterial Activity

The antibacterial activity of mangrove leaf extract was performed with disc diffusion method. Whatman filter paper discs (5mm) were impregnated with various concentrations (50, 100, 150, 200, 250 & 300 $\mu\text{g/ml}$) of leaf extract. The inoculated plates were incubated for 24 hr at room temperature and the inhibition zones around the discs were measured. All the results were expressed with mean \pm standard error.

2.6 Minimum Inhibitory Concentrations

Minimal Inhibition Concentration of mangrove leaf extract was determined in 5 concentrations (50 -300 $\mu\text{g/ml}$) with blank (extract in Muller Hinton broth). The inoculated bacteria in test tubes are incubated for 24hr in ambient temperature. The results are noted as well growth (+) and inhibited (-). Tetracycline (1 mg/ml) was used as standard and DMSO was used as negative control.

3. RESULTS

On measuring the zone of inhibitions for each pathogen at different concentrations ranging from 50 $\mu\text{g/ml}$ to 300 $\mu\text{g/ml}$. The zone of inhibition for MDRSA was minimum i.e. 2mm at 50 $\mu\text{g/ml}$ of concentration and maximum i.e. 13mm at 30 $\mu\text{g/ml}$ of concentration. The zone of inhibition for MRSA as well as VRSA was minimum i.e. 2mm at 100 $\mu\text{g/ml}$ of concentration and maximum i.e. 13mm and 9mm respectively at 300 $\mu\text{g/ml}$ of concentration (Table 1). The zone of inhibition at 300 $\mu\text{g/ml}$ of concentration for MDRSA and MRSA showed higher effect (13mm) when compared to VRSA (9mm). The minimum inhibitory concentration for MDRSA was 250 $\mu\text{g/ml}$ of concentration, MRSA was 200 $\mu\text{g/ml}$ of concentration and VRSA was $\mu\text{g/ml}$ of 150 $\mu\text{g/ml}$ of concentration (Table 2).




Fig. 2. Images showing antibacterial activity of *Avicennia marina* extract against (A) MRSA, (B) VRSA, (C) MDRSA

Table 1. Table showing zone of inhibition produced by *Avicennia marina* on MDRSA, MRSA and VRSA . The values are given in triplicate with mean \pm SE.

Concentration of <i>Avicennia marina</i> ($\mu\text{g/ml}$)	MDRSA	MRSA	VRSA
0	0	0	0
50	2 \pm 0.86	0	0
100	5 \pm 1.3	2 \pm 0.57	2 \pm 0.8
150	8 \pm 1.07	5 \pm 1.2	4 \pm 1.07
200	11 \pm 1.2	6 \pm 1.4	5 \pm 1.2
250	12 \pm 1.4	9 \pm 1.2	8 \pm 1.4
300	13 \pm 1.2	13 \pm 1.4	9 \pm 1.2

Table 2. Table showing MIC for MDRSA, MRSA and VRSA

$\mu\text{g/ml}$	50	100	150	200	250	300	MIC ($\mu\text{g/ml}$)
MDRSA	+	+	+	+	-	-	250
Tetracycline	+	+	-	-	-	-	150
MRSA	+	+	+	-	-	-	200
Tetracycline	+	+	-	-	-	-	150
VRSA	+	+	+	+	-	-	250
Tetracycline	+	+	-	-	-	-	150

4. DISCUSSION

The heartwood of Red sanders is known to have antipyretic, anti-inflammatory, anthelmintic, tonic, hemorrhage, dysentery, aphrodisiac, and diaphoretic activities. It has also been used as a cooling agent. Ethanol extract of stem bark was reported to possess anti-hyperglycaemic activity. The wood in combination with other drugs is also prescribed for snake bites and scorpion stings. Phytochemical investigations of aqueous and ethanol extracts of stem bark has revealed the presence of alkaloids, phenols, saponins, glycosides, flavonoids, triterpenoids, sterols, and tannins [3,32,33].

The present interest in medicinal plants as therapeutic agents has progressed in different

parts of the world because of the increasing incidence of bacterial resistance to chemical drugs and the emergence of new pathogenic bacterial strains. On carrying out in vitro testing of a large number of plants against various bacterial strains, it has been demonstrated that extracts and pure compounds of numerous medicinal plants are very effective against bacterial strains [34].

In India, the epidemiology of MRSA has been changing over the past few decades. The resistance of MRSA to B-lactam like penicillin and amoxicillin is 100% [35]. Previous studies have revealed that plant extracts exhibit antibacterial activity against pathogenic bacteria stains such as, *Staph. aureus*, *E.coli*, *Pseudomonas* and antibiotic resistant strains. In

a similar study conducted by Dhayanithi et al., 2012 reported that *A. marina* extract showed inhibitory activity against multi-drug resistant *S. aureus*. *Staphylococcus aureus* strains showed resistance to ciprofloxacin, methicillin and vancomycin based on the antibiotic susceptibility test [36].

Human oral cavity is normally sterile at birth but soon it possesses a predominant streptococcal microbiota. The species often related to "viridans streptococci" comprises *Streptococcus salivarius*, *Streptococcus sanguis*, *Streptococcus mitis*, and *Streptococcus mutans* [37]. Dental caries, a microbial disease is caused by α -hemolytic streptococci usually referred to as *S. viridans*, a major component of dental plaque and primarily responsible for pediatric dental health related problems. [38]

In a study conducted by Mahasneh 2002 using *Avicennia marina* leaves extracts in methanol showed antibacterial activity against *B. subtilis*, *S. aureus*, and *Streptococcus pyogenes* with approximately 20 mm zone of inhibition. Butanol crude extract of the plant exhibited a potent antibacterial activity against both Gram-positive and Gram-negative bacteria including *E. coli*, *B. cereus*, *S. aureus*, *P. aeruginosa*, *A. flavus*, *C. albicans* with zone of inhibition measuring 15 mm, 14 mm, 10 mm, 14 mm, 13 mm, and 14 mm, respectively [39] Many research were carried out translated to publication [40-42] recently by our team.

A study conducted by Okla et al., 2021 has shown that root and leaf extracts of *A. marina* exhibited antibacterial activity. Ethanol root extract proved to be effective against bacterial strains, *P. aeruginosa*, *B. subtilis*, *S. aureus*, and *E. coli* as well as fungal strains *Aspergillus fumigatus* and *Candida albicans*. Leaf extract in ethyl acetate displayed significant antibacterial activity against *S. aureus* and *E. coli*.

5. CONCLUSION

The present study showed effective antibacterial activity of *Avicennia marina* against MDRSA, MRSA, VRSA. Hence, this extract may be used for infections against resistant *Staphylococcus aureus*.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The study was ethically approved by the Scientific Review Board, Saveetha Dental College, Chennai.

ACKNOWLEDGEMENT

The authors thank the management of Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences for all support to carry out this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Devaraj E, Roy A, Veeraragavan GR, Magesh A, Sreeba AV, Arivarasu L, et al. β -Sitosterol attenuates carbon tetrachloride-induced oxidative stress and chronic liver injury in rats [Internet]. Naunyn-Schmiedeberg's Archives of Pharmacology. 2020;393:1067–75. Available: <http://dx.doi.org/10.1007/s00210-020-01810-8>
2. Anitha R, Ashwini S. Antihyperglycemic activity of *Caralluma fimbriata*: An In vitro approach [Internet]. Pharmacognosy Magazine. 2017;13:499. Available: http://dx.doi.org/10.4103/pm.pm_59_17
3. Anitha R, Prathoshni S, Lakshmi T. The effect of capsicum oleoresin on nitric oxide production and nitric oxide synthase gene expression in macrophage cell line [Internet]. Pharmacognosy Research. 2018;10:343. Available: http://dx.doi.org/10.4103/pr.pr_46_18
4. Cinthura C, Thangavelu L, Rajeshkumar S, Gurunadhan D, Pradeep Kumar R, Roy A. COX2 Inhibitory activity of *Abutilon indicum*-An Invitro Study [Internet]. Indian Journal of Public Health Research & Development. 2019;10:3523. Available: <http://dx.doi.org/10.5958/0976-5506.2019.04131.7>
5. Karthikeyan G, Lakshmi T, Rajeshkumar S, Roy A, Gurunadhan D, Geetha RV, et al. Glucose Uptake Potential in L6 Myotubes by *Ficus racemosa* [Internet]. Indian Journal of Public Health Research & Development. 2019;10:3527.

- Available: <http://dx.doi.org/10.5958/0976-5506.2019.04132.9>
6. Suhasini SJ, Jennifer Suhasini S, Roy A, Sosa G, Lakshmi T. The Cytotoxic effect of *Caralluma fimbriata* on KB cell lines [Internet]. *Research Journal of Pharmacy and Technology*. 2019;12:49-95. Available: <http://dx.doi.org/10.5958/0974-360x.2019.00865.5>
 7. Roy A, Rasheed A, Sleeba AV, Rajagopal P. Molecular docking analysis of capsaicin with apoptotic proteins. *Bioinformation*. 2020 Jul 31;16(7):555–60.
 8. Aneesa NN, Anitha R, Varghese S. Antidiabetic Activity of Ajwain Oil in Different Models. *J Pharm Bioallied Sci*. 2019 Apr;11(2):142–7.
 9. Krishna RN, Anitha R, Ezhilarasan D. Aqueous extract of fruit pulp exhibits antihyperglycaemic activity. *Avicenna J Phytomed*. 2020 Sep;10(5):440–7.
 10. Meenapriya M, Anitha R, Lakshmi T. Effect of Lutein on Cytochrome P450 (Isoform CYP3A4) - An in vitro Study [Internet]. Vol. 10, *Pharmacognosy Journal*. 2018. p. 1093–5. Available:<http://dx.doi.org/10.5530/pj.2018.6.185>
 11. Turner M. Ecology: Mangrove maintenance. *Nature*. 2015;526(7574):515.
 12. Agoramoorthy G, Chandrasekaran M, Venkatesalu V, Hsu MJ. Antibacterial and antifungal activities of fatty acid methyl esters of the blind-your-eye mangrove from India [Internet]. *Brazilian Journal of Microbiology*. 2007;38:739–42. Available: <http://dx.doi.org/10.1590/s1517-83822007000400028>
 13. Bandaranayake WM. Wetlands Ecology and Management [Internet]. Vol. 10. 2002. p. 421–52. Available from: <http://dx.doi.org/10.1023/a:1021397624349>
 14. Bandaranayake WM. Economic, Traditional and Medicinal Uses of Mangroves. 1999. 82 p.
 15. Hogarth PJ. Mangroves and Seagrasses [Internet]. *The Biology of Mangroves and Seagrasses*. 2015:1–7. Available:<http://dx.doi.org/10.1093/acprof:oso/9780198716549.003.0001>
 16. Marine Enzymes Biotechnology: Production and Industrial Applications, Part III - Application of Marine Enzymes. Academic Press; 2017:228.
 17. Rajaram R, Ganeshkumar A, Muralisankar T, Sivaperumal P. Bioaccumulation of metals in mangroves and salt marshes collected from Tuticorin coast of Gulf of Mannar marine biosphere reserve, Southeastern India. *Mar Pollut Bull*. 2020;160:111599.
 18. Kamala K, Sivaperumal P, Thilagaraj R, Natarajan E. Bioremediation of Sr² ion radionuclide by using marine *Streptomyces* sp. CuOff24 extracellular polymeric substances [Internet]. *Journal of Chemical Technology & Biotechnology*; 2019. Available:<http://dx.doi.org/10.1002/jctb.6018>
 19. Sivaperumal P, Kamala K, Rajaram R. Biosorption of Long Half-life Radionuclide of Strontium Ion (Sr) by Marine Actinobacterium *Nocardiostrictum* sp. 13H [Internet]. *Geomicrobiology Journal*. 2018; 35:300–10. Available:<http://dx.doi.org/10.1080/01490451.2017.1350891>
 20. Bandaranayake WM. Mangroves and Salt Marshes [Internet]. 1998;2:133–48. Available:<http://dx.doi.org/10.1023/a:1009988607044>
 21. Devi VS, Subathra Devi V, Gnanavel BK. Properties of Concrete Manufactured Using Steel Slag [Internet]. *Procedia Engineering*. 2014;97:95–104. Available:<http://dx.doi.org/10.1016/j.proeng.2014.12.229>
 22. Gupta P, Ariga P, Deogade SC. Effect of Monopoly-coating Agent on the Surface Roughness of a Tissue Conditioner Subjected to Cleansing and Disinfection: A Contact Profilometric Study. *Contemp Clin Dent*. 2018;(Suppl 1):S122–6.
 23. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog*. 2018;117:68–72.
 24. Needhidasan S, Samuel M, Chidambaram R. Electronic waste - an emerging threat to the environment of urban India. *J Environ Health Sci Eng*. 2014;12(1):36.
 25. Rasoanaivo P, Petitjean A, Ratsimamanga-Urverg S, Rakoto-Ratsimamanga A. Medicinal plants used to treat malaria in Madagascar. *J Ethnopharmacol*. 1992;37(2):117–27.
 26. Rajeshkumar S, Kumar SV, Ramaiah A, Agarwal H, Lakshmi T, Roopan SM. Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and

- evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. *Enzyme Microb Technol.* 2018 Oct;117:91–5.
27. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation. *Biocatal Agric Biotechnol.* 2019 May 1;19:101138.
 28. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.–mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res.* 2020 Mar 1;27(8):8166–75.
 29. Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using *Gymnema sylvestris* leaf extract and evaluation of its antibacterial activity [Internet]. *South African Journal of Chemical Engineering.* 2020;32:1–4.
Available:<http://dx.doi.org/10.1016/j.sajce.2019.11.005>
 30. Rajasekaran S, Damodharan D, Gopal K, Rajesh Kumar B, De Pours MV. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel.* 2020 Oct 1;277:118166.
 31. Veerasamy R, Roy A, Karunakaran R, Rajak H. Structure-Activity Relationship Analysis of Benzimidazoles as Emerging Anti-Inflammatory Agents: An Overview. *Pharmaceuticals* [Internet]. 2021 Jul 11;14(7).
Available:<http://dx.doi.org/10.3390/ph14070663>
 32. Dhayanithi J, Rajeshkumar S, Roy A, Lakshmi T. Preparation and Evaluation of Antifungal Activity of Arrow Root Mediated Selenium Nanoparticles Against *Candida Albicans* -. *Journal of Complementary Medicine Research.* 2020;11(5):83–8.
 33. Role of Nanomedicine in Novel Corona Virus Pandemic: A perspective [Internet]. 2020
[Cited 2021 Aug 31].
Available:<http://bbrc.in/bbrc/role-of-nanomedicine-in-novel-corona-virus-pandemic-a-perspective/>
 34. Mahady GB, Huang Y, Doyle BJ, Locklear T. Natural Products As Antibacterial Agents [Internet]. *Bioactive Natural Products (Part O).* 2008:423–44.
Available: [http://dx.doi.org/10.1016/s1572-5995\(08\)80011-7](http://dx.doi.org/10.1016/s1572-5995(08)80011-7)
 35. Vidhani S, Mehndiratta PL, Mathur MD. Study of methicillin resistant *S. aureus* (MRSA) isolates from high risk patients. *Indian J Med Microbiol.* 2001 Apr;19(2):13–6.
 36. Dhayanithi NB, Ajith Kumar TT, Ganesha Murthy R, Kathiresan K. Isolation of antibacterials from the mangrove, *Avicennia marina* and their activity against multi drug resistant *Staphylococcus aureus* [Internet]. *Asian Pacific Journal of Tropical Biomedicine.* 2012;2:S1892–5.
Available: [http://dx.doi.org/10.1016/s2221-1691\(12\)60516-4](http://dx.doi.org/10.1016/s2221-1691(12)60516-4)
 37. Pearce C, Bowden GH, Evans M, Fitzsimmons SP, Johnson J, Sheridan MJ, et al. Identification of pioneer viridans streptococci in the oral cavity of human neonates. *J Med Microbiol.* 1995 Jan;42(1):67–72.
 38. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the normal bacterial flora of the oral cavity. *J Clin Microbiol.* 2005 Nov;43(11):5721–32.
 39. Mahasneh AM. Screening of some indigenous Qatari medicinal plants for antimicrobial activity [Internet]. *Phytotherapy Research.* 2002;16: 751–3.
Available:<http://dx.doi.org/10.1002/ptr.1037>
 40. Okla MK, Alatar AA, Al-amri SS, Soufan WH, Ahmad A, Abdel-Maksoud MA. Antibacterial and Antifungal Activity of the Extracts of Different Parts of *Avicennia marina* (Forssk.) Vierh [Internet]. *Plants.* 2021;10:252.
Available:<http://dx.doi.org/10.3390/plants10020252>
 41. Rajendran R, Kunjusankaran RN, Sandhya R, Anilkumar A, Santhosh R, Patil SR. Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An in Vitro Study. *Pesqui Bras Odontopediatria Clin Integr.* 2019; 19(0):4668.

42. Kumar MS, Vamsi G, Sripriya R, Sehgal PK. Expression of matrix metalloproteinases (MMP-8 and -9) in chronic periodontitis patients with and without diabetes mellitus. J Periodontol. 2006;77(11):1803–8.

© 2021 Ramasubramanian et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77720>