



# In Planta Androgenesis (Paternal Origin) in *Saccharum X Erianthus* Intergeneric Hybrid

K. V. Sushir <sup>a++</sup>, Apuunu C. <sup>b++</sup> and B. S. Thorat <sup>c#\*</sup>

<sup>a</sup> V.S.I., Pune, India.

<sup>b</sup> S.B.I., Coimbatore, India.

<sup>c</sup> Dr. B.S.K.K.V., Dapoli, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors contributed equally to this work. Authors KVS, AC and BST wrote the manuscript effectively. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i91305>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121865>

Original Research Article

Received: 20/06/2024

Accepted: 22/08/2024

Published: 28/08/2024

## ABSTRACT

The present investigation was carried in crosses made between *Saccharum species* hybrid clones i.e. Co 419, Co 7201 and Co 7224 (2n = 108-112) and *Erianthus arundinaceus* viz. IK 76-91 (2n = 60) and IK 76-99 (2n = 60) for introducing better ratoon ability, vigour, wide ecological adaptability, extensive root system, drought and water logging resistance, high biomass and disease resistance into modern sugarcane cultivars during 2020-23 at VSI-Sugarcane Breeding Centre, Amboli. Parental apomixis (paternal origin) has been recently noticed and it is unique in *Saccharum X Erianthus* intergeneric cross seedlings. A total of fifty-one seedlings were obtained from above

<sup>++</sup>Senior Scientist;

<sup>#</sup>Assistant Seed Production Officer;

<sup>\*</sup>Corresponding author: E-mail: [balajithorat2014@gmail.com](mailto:balajithorat2014@gmail.com);

**Cite as:** Sushir, K. V., Apuunu C., and B. S. Thorat. 2024. "In Planta Androgenesis (Paternal Origin) in *Saccharum X Erianthus* Intergeneric Hybrid". *Journal of Advances in Biology & Biotechnology* 27 (9):355-61. <https://doi.org/10.9734/jabb/2024/v27i91305>.

three intergeneric crosses and out of these eighteen seedlings were reported as androgenetic seedlings as the female act as a surrogate mother. At the early stages, seed germination of seedlings was found two types i.e., one type resembling *Saccharum* and other resembling *Erianthus* with deep purple coloration. Morphologically androgenetic seedlings were similar to the male parents IK 76-91 and IK 76-99 with tall large tufted grass like clumps, vegetative stem with thick reproductive buds, root zone narrow and only one row of root eyes, the lamina gradually passes into leaf sheath with no auricle and dewlaps. Cytological studies were made in these seedlings and during metaphase the somatic chromosome number was observed to be  $2n = 60$  rather than expected  $2n = 84-90$ . This paternal origin may be due to the ability of *Erianthus* male gametes to produce an embryo within the seed tissues of *Saccharum species* hybrids but without genetic contribution of the seed. Thus, *in planta* androgenesis is achieved through the combination of embryogenic behavior of male gametes which has entered in the ovule and the ability of ovules to act as a surrogate mother. They could derive either from the fusion of two male gametes or from the early natural diploidization of the haploid embryo. Such homozygous true breeding seedlings of *Saccharum X Erianthus* with hybrid vigour are extremely important in crop improvement. This system provides an unparalleled opportunity to shorten the breeding cycle and fix agronomic traits in the homozygous state. These androgenetic plants with *Saccharum* cytoplasm and *Erianthus* genome are male sterile making them highly useful for further hybridization and sugarcane improvement programme.

**Keywords:** Androgenesis; cytology; *Erianthus*; intergeneric hybrid and *Saccharum*.

## 1. INTRODUCTION

Modern sugarcane cultivars are complex interspecific hybrids ( $2n = 100-130$ ). They were developed from interspecific crosses made early this century among relatively few clones of *Saccharum officinarum* ( $2n = 80$ ), the sugar producing species and *S. spontaneum* L. ( $2n = 40-128$ ) a wild species. Breeders are concerned by the potentially narrow genetic base of contemporary commercial crops. The genera *Saccharum* L., *Erianthus* Mychx. Section *Ripidium* Henrad, *Miscanthus* Anderss, *Sclerostachya* (Hack.) A. Camus and *Narenga* Bor., which are closely related, were considered to be part of 'Saccharum complex' [1,2] sharing a common gene pool.

*Erianthus arundinaceus* (Retz.) Jeswiet ( $2n = 60$ ) belongs to the genus *Erianthus* sect. *Ripidium* is included in 'Saccharum complex'. It is a large grass with tall and thick stalk resembling sugarcane and has a great potential as a contributor of germplasm to current cultivars for better ratoon ability, vigour, wide ecological adaptability, extensive root system, drought and water logging resistance, high biomass and disease resistance [3,4].

The genetic base of sugarcane is considered to be very narrow and gene introgression studies are being done to transfer the genes from wild species or related genera. Numerous attempts have been made to cross *E. arundinaceus* with

sugarcane to introduce these characters into modern cultivars [5,6]. However, no conclusive results have been achieved to date. Indeed, in only a few cases have conclusive evidences of true intergeneric hybrids being documented [7,4,8]. One of the major obstacles in the past has been the identification of true hybrids using morphological characters.

Intergeneric crosses involving *Saccharum* spp hybrid as a female were made with *E. arundinaceus* as a male at Distant Hybridization Garden, Agali (Kerala), India in 2008 and morphological and cytological studies were taken up in the fifty-one seedlings obtained from the *Saccharum X Erianthus* three different crosses (ICAR) [9]. Hence, this study was undertaken for to shorten the breeding cycle by developing male sterile lines and fix agronomic traits in the homozygous state.

## 2. MATERIALS AND METHODS

The *Saccharum* spp. hybrid Co 419, Co 7201 and Co 7224 were pollinated with *E. arundinaceus* (Retz.) Jeswiet clone IK 76-91 and IK 76-99 and the seedlings raised were maintained in the field during 2021-23 at VSI-Sugarcane Breeding Centre, Amboli. The somatic chromosome of the parental clones and the seedlings were determined by root tip squash technique. Actively growing root tips from cane cutting planted in pots with river sand were excised and treated in a saturated solution of  $\alpha$ -

bromonaphthalene for about two hours. After thoroughly washing in distilled water the root tips were fixed in Ostergren and Heneen's [10] fixative for about 24 hours. The root tips were hydrolyzed in 1N HCL for 15 minutes at  $58 \pm 2$  °C and stained in Leuco basic fuchsin for 30 minutes and squashed in 1% acetocarmine. Well spread mitotic chromosomes were counted and photographed. The pollen fertility of the parents and the seedlings were tested with the help of Differentials stain [11]. The red stained pollen is categorised as the fertile pollen and unstained pollen as sterile.

### 3. RESULTS AND DISCUSSION

Nine seedlings were obtained from *Saccharum* species hybrid; Co 7224 florets pollinated with *E. arundinaceus* clone IK 76-99 pollen, whereas six seedlings from the Co 419 X IK 76-91 and thirty-six seedlings from the Co 7201 X IK 76-91, respectively. All the seedlings in general were fast growing with good vigour. Among these fifty-one seedlings from the three crosses, eighteen seedlings were found to be androgenetic seedlings (Co 419 X IK 76-91: 06; Co 7224 X IK 76-99: 08 and Co 7201 X IK 76-91: 04). Morphologically these androgenetic seedlings were similar to the male parents IK 76-91 and IK 76-99 with tall large tufted grass like clumps, vegetative stem with thick reproductive buds, root zone narrow and only one row of root eyes, the lamina gradually passes into leaf sheath with no auricle and dewlaps. The chromosome number of *Saccharum* species hybrid Co 419, Co 7201 and Co 7224 and *E. arundinaceus* clone IK 76-99 were determined to be  $2n = 108-112$  and  $2n = 60$  respectively. Intergeneric hybrid between these had expected  $n + n$  transmission with  $2n = 84-90$ . During cytological study of seedlings about eighteen seedlings from the three different intergeneric crosses, the somatic chromosome number at metaphase recorded with  $2n=60$  (Fig. 1b) and all the other seedlings were  $n + n$  hybrids with chromosome number with  $2n= 84$  to 90.

These eighteen intergeneric cross seedlings were morphologically and cytologically identical with the *Erianthus* male parent and recorded as androgenetic plant. No hairs on the eye bud (Fig. 1g, 1j), only one row of the root eyes (Fig. 1h, 1k) and ligules are reduced (Fig. 1i, 1l) were present in all the androgenetic seedlings like male parents IK 76-91 and IK 76-99 (Fig. 1n, 1o). Two instances were recorded in the androgenetic seedlings, it had the similarity in morphologically

and cytologically to the donor male parents. In looking to the morphological features of the *Saccharum* X *Erianthus* hybrid seedlings, the expected cytoplasm from *Saccharum* was missing with an element of uncertainty and it could derive either from the fusion of two male gametes or from the early natural diploidization of the haploid embryo. The pollen fertility was also studied in the parent and hybrid seedlings. The androgenetic seedlings (Fig. 1n, 1o) although had morphologically and cytologically similar to male parent but these seedlings had recorded with 100% sterile i.e. unstained pollen (Fig. 1p). The female parent Co 7224 recorded with 28% fertility and showed both the red and unstained pollen (Fig. 1q) and the male parents IK 76-91 and IK 76-99 were 100% pollen fertile with complete red pollen (Fig. 1r). Until recently, apomixis had been reported only in Angiosperms. The highly endangered Mediterranean conifer, *Cupressus dupreziana* A. Camus (or Tassil cypress), is to our knowledge the first plant species where in planta "parental apomixis" (i.e. embryogenic development of diploid pollen in seed tissues) was hypothesized [12].

Sexual reproduction is the general rule in higher plants and animals. More than 99% of multicellular eukaryotes reproduce sexually [13,14,15]. The sexual cycle follows the alteration of haploid and diploid cells delimited by meiosis and syngamy, two events that modify the nuclear chromosome number [16]. However, in some asexual species in 40 angiosperm families (mainly Poaceae, Asteraceae, Rosaceae and Rutaceae) exhibit pattern of apomixis where the embryo is derived solely from maternal tissues, either somatic cell or a gamete with the somatic chromosome number (unreduced gamete) [17]. In rare cases, the parthenogenetic development of a reduced gametophyte produces a haploid embryo [18]. This last feature, also called non-recurrent apomixis, occurs naturally in some female gametophytes. It is also intensively used in breeding programs through the in vitro culture of male gametophytes (androgenesis). Diploid homozygous genotypes can then be obtained through or artificial doubling of the chromosome number.


The *E. arundinaceus* clones IK 76-91 and IK 76-99 were collected from Kalimantan province in Indonesia and is with chromosome number  $2n = 60$  as that in other clones of this species collected from this region. They are hexaploids with basic chromosome number  $X = 10$  [19].

Sreenivasan [20] reported that *Saccharum X Erianthus* hybridization looked most promising for transferring many desirable characters to present day commercial varieties, since these hybrids do not show significant reduction in sucrose content especially when *E arundinaceus* is used as a male parent. In the present investigation, the intergeneric hybrid seedlings recorded with  $2n = 60$  chromosome number which was not expected and it was to be  $2n = 84-90$  with normal  $n + n$  chromosome transmission. These hybrids were identical to the male parent IK 96-91 and IK 76-99 not only in respect of chromosome constitution but also in morphological features. Generally, in the intergeneric hybrid between *Saccharum X Erianthus*, the contribution of cytoplasm could be of *Saccharum* and the nuclear genome from both the genera chromosome contribution. In this study, eighteen hybrid seedlings from three different crosses, the *Saccharum* cytoplasm was eliminated by the *Erianthus* genera and occupied both the cytoplasm and nuclear genome and recorded as androgenetic plant. When *Erianthus* was used as a male parent in hybridization with *S. officinarum* and Co 7201, two instances of androgenesis were observed. The seedlings had the morphology and chromosome number of the male parent [21].

The peculiarity that is almost unique in sugarcane hybridization is that no prediction can be made as to what chromosome number is a particular cross would be like. Many introgression programmes have been attempted without to success to introduce characters from *E. arundinaceus* to *Saccharum species* hybrids. Three major difficulties occurred at different steps of these intergeneric crosses programme [22]. The first difficulty is the production of progeny from intergeneric crosses with *E. arundinaceus*. Lee [23] showed that incongruity between the pollen and pistils of the two genera occurs since he observed that the *Erianthus* pollen tube often did not reach the micropyle of the *Saccharum* ovules. They either stopped growing toward it or they lost directional control. This uncoordinated pollination process happens between distinct species [24] and resulted in limited intergeneric fecundation. In addition to sugarcane flowers are hermaphroditic with varying degrees of male sterility clones with less than 30% viable pollens are used as females in crossing. Thus because of incongruity between the pollen and pistils, even if the *Erianthus* parent is a strong male and female parent has little

fertile pollen, the few crosses that produce seed generate mostly self-pollinated individual. The second difficulty was the identification of genuine hybrids using morphological characters, which has been highlighted by use of molecular diagnostic tools [4]. The third difficulty is to maintain the genuine hybrids, which often are very weak.

Using cytological tools, we were able to analyse chromosome number of eighteen intergeneric hybrid seedlings with  $2n = 60$  which were identical to the male *Erianthus* parent IK 76-91 and IK 76-99 and morphologically these plants were also similar in all respect with male parent. The cytoplasm was expected from *Saccharum* genera as it was used as a female parent. This paternal origin may be due to the ability of *Erianthus* pollen to produce an embryo within the seed tissues of *Saccharum* species but without genetic contribution of the seed, as reported Pichot et al. [25] when *Cupressus sempervirens* seed trees were pollinated by *C. dupreziana* pollen. The androgenetic process occurred from unreduced diploid pollen of *C. dupreziana* and all the progeny from *C. sempervirens* seed trees were diploid. They could derive either from the fusion of genetically identical two male gametes (i.e. deriving from the same pollen) or from the early natural diploidization of the haploid embryo. Although the production of all maternal progeny by apomixis or parthenogenesis is a rather frequent phenomenon, the production of all paternal progeny has been very rarely reported [26]. To our knowledge, *C. dupreziana* is the only plant in which progeny are produced by the apomictic development of pollen grains. The scarcity of androgenetic cases reported compared to the amount of gynogenic case reports may be explained not only by the probable evolutionary dead end of this reproductive process but also by the difficulty in detecting it [26]. The male component ability to produce an embryo without female gamete contribution may not be so rare. In fact, it is intensively used in Angiosperms to produce haploid genotypes from in vitro culture of immature anthers. Thus, favourable growth conditions combined with a lack of syngamy opportunity may often lead to androgenesis. This paternal origin may be due to the ability of *Erianthus* male gametes to produce an embryo within the seed tissues of *Saccharum species* hybrids but without genetic contribution of the seed.



**Fig. 1. Somatic chromosome number at prophase, metaphase and anaphase (Fig. 1a, 1b and 1c), No hairs on eye bud (Fig. 1g, 1j), only one row of the root eyes (Fig. 1h, 1k) and reduced ligules (Fig. 1i, 1l), Androgenetic seedlings (Fig. 1n, 1o), unstained pollen grains (Fig. 1p) found in androgenic seedlings, stained & unstained pollen grains (Fig. 1q) found in female clones, and fully stained pollen grains (Fig. 1r) found in male clones, morphological traits hair on eye bud (Fig. 1d), 2-3 rows of root eyes (Fig. 1e) and hairs on ligules (Fig. 1f)**

#### 4. CONCLUSION

It is concluded that *in planta* androgenesis is achieved through the combination of embryogenic behaviour of male gametes which has entered in the ovule and the ability of ovules to act as a surrogate mother. They could derive either from the fusion of two male gametes or from the early natural diploidization of the haploid embryo. This system provides an unparalleled opportunity to shorten the breeding cycle and fix agronomic traits in the homozygous state. Such homozygous true breeding androgenetic seedlings of *Saccharum X Erianthus* with hybrid vigour are extremely important for further sugarcane crop improvement.

#### CONFERENCE DISCLAIMER

Abstract part of this manuscript was previously presented and published in the conference: 10Th GERMPLOSM & BREEDING 7Th MOLECULAR BIOLOGY WORKSHOP, MACEIÓ-BRAZIL / 15-20 MAY 2011, Web Link of the proceeding: [https://www.comunicventos.com.br/\\_HOTSITES/germplasm/\\_ARQUIVOS/ABSTRACT.pdf](https://www.comunicventos.com.br/_HOTSITES/germplasm/_ARQUIVOS/ABSTRACT.pdf)

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### ACKNOWLEDGEMENT

The authors are thankful to Dr. T. V. Sreenivasan Ex. Director, SBI, Coimbatore and Hon. Mr. Sambhaji Kadu Patil, Director General, VSI for their constant encouragement.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Mukherjee SK. Origin and distribution of *Saccharum*, Bot. Gaz. 1957;119:55-61.
2. Daniels J, Smith P, Paton N, Williams CA. The origin of the genus *Saccharum*. Sugarcane Breed. Newsl. 1975;36:24-39.

3. Roach BT. A programme for sugarcane improvement from genetic diversity: background and preliminary results, Proc. internal Soc. Sugarcane Technol. 1989; 20:900-909.
4. D'. Hont A, Rao PS, Feldman P, Rivet L, Islam-Faridi N, Taylor P, Glazmann JC. Identification and characterization of sugarcane intergeneric hybrid, *Saccharum officinarum X Erianthus arundinaceus* with molecular markers and DNA in situ hybridization. Theor. Appl. Genet. 1995;91: 320-326.
5. Hale B, Ferrie AMR, Chellamma S, Samuel JP, Phillips GC. Rogeneration-based double Haploidy: Past, Present and Future Perspectives, Front. Plant. Sci. 2022;12: 751230.
6. Samantaray, et al. High-frequency androgenic green plant regeneration in Indica Rice for Accelerated Breeding, Springer, Cham. 2020;1.
7. Nagai C, Ahloowalia BS, Tew TL. Somaclonal variants from an intergeneric hybrid: *Saccharum* spp. hybrid X *Erianthus arundinaceus*. Euphytica. 1991;53:193-199.
8. Besse P, McIntyre CL, Burner DM, de Almeida CG. Using genomic slot blot hybridization to assess intergeneric *Saacharum X Erianthus* hybrids (Andropogoneae-Saccharinae). Genome. 1997a;40:428-432.
9. ICAR. Available:<https://sugarcane.icar.gov.in/index.php/genetic-studies/>
10. Ostergreen G, Heneen K. A quash technique for chromosome morphological studies. Hereditas. 1962;48:332-342.
11. Alexander MP. Differential staining of aborted and non-aborted pollen. Stain Technology. 1969;44:117-122. Available:<https://doi.org/10.3109/10520296909063335>
12. Pichot C, Fady B, Hochu I. Lack of mother tree alleles in zymograms of *Cupressus dupreziana* A Camus emryo. Ann. For. Sci. 2000;57:17-22.
13. Bell G. The masterpiece of Nature: The evolution and Genetics of Sexuality. University of California Press, Berkeley, CA; 1982.
14. Barton NH, Charlesworth B. Why sex and recombination? Science. 1998;281:1986-1990.
15. West SA, Lively CM, Read AF. A pluralistic approach to sex and

- recombination. J. Evol. Biol. 1999;12:1003-1012.
16. Vrijenhoek RC. Clonal organism and benefits of sex. Adv. Mol. Ecol. 1998; 306:151-172.
  17. Vielle Calzada JP, Crane CF, Stelly DM. Apomixis: The asexual revolution. Science. 1996;274:1322-1323.
  18. Foroughi-Wehr B, Wenzel G. Ro and parthenogenesis in Plant Breeding: Principles and Prospects, edited by M.D. Hayward, N.O. Bosemark and I. Romagosa Chapman and Hall, London. 1993;261-277.
  19. Besse P, McIntyre C. L. 1998. Chromosome in situ hybridization of ribosomal DNA of *Erianthus sect. Ripidium species* with varying chromosome numbers confirms  $x = 10$  in *Erianthus sect. Ripidium*. Genome, 42: 270-273.
  20. Sreenivasan TV, Ahloowalia BS, Heinz DJ. Cytogenetics, In. D. J. Heinz (ed). Sugarcane Improvement through Breeding Elsevier, Netherland; 1987.
  21. Anon. (T. V. Sreenivasan and N. C. Jalaja) Annual Report. Sugarcane Breeding Institute Coimbatore. 1989-90;26.
  22. Piperidis G, Christopher MJ, Carol BJ, Berding N, D'Hont A. Molecular contribution to selection of intergeneric hybrids between sugarcane and the wild species *Erianthus arundinaceus*. Genome. 2000;43:1033-1037.
  23. Lee DJ. Enhancement of *Saccharum* spp. hybrid material by introgression with *Erianthus arundinaceus* germplasm. Ph.D. Thesis, James Cook University of North Queensland, Townsville, Queensland; 1995.
  24. Hogenboom N. Incompatibility and incongruity: Two different mechanisms for the non-functioning of intimate partner relationship. Proc. R. Soc. Lond. Biol. Sci. 1975;188:361-375.
  25. Pichot C, El Maataoui M, Raddi S, Raddi P. Surrogate mother for endangered Cupressus. Nature. 2001;412:39.
  26. McKone MJ, Halpern SL. The evolution of androgenesis Ame. Nat. 2003;161:641-656.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121865>