



# Brunner's Gland Hamartoma of the Duodenum as a Cause of Intestinal Obstruction in a Young Patient: A Surgical Approach

Imad Laamri <sup>a\*</sup>, Mohammed Dady <sup>a</sup>, Sabrillah Echiguer <sup>a</sup>,  
Yousra Ouhammou <sup>a</sup>, Oumaima Lahnaoui <sup>a</sup>,  
Amine Souadka <sup>a</sup>, Mohammed Anass Majbar <sup>a</sup>  
and Amine Benkabbou <sup>a</sup>

<sup>a</sup> Surgical Oncology Department, National Institute of Oncology, University Mohammed V in Rabat, Rabat, Morocco.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123210>

Case Report

Received: 26/07/2024

Accepted: 28/09/2024

Published: 03/10/2024

## ABSTRACT

The Brunner's gland hamartoma is a benign tumor of the duodenum, but it has malignant potential with a very low risk of progressing to adenocarcinoma. It is rare, accounting for less than 1% of primary tumors of the small intestine. Furthermore, its clinical manifestations are nonspecific, its etiology is poorly understood, and the treatment strategy needs to be tailored accordingly. We

\*Corresponding author: E-mail: laamriimad@gmail.com;

report a case of a patient with a Brunner's gland hamartoma of the duodenum who presented with intestinal obstruction and was treated with surgical resection of the polyp via duodenotomy.

**Keywords:** Brunner's gland; duodenum; hamartoma; hyperplasia.

## 1. INTRODUCTION

"The Brunner's glands, described by the anatomist Brunner in 1688, are submucosal mucous glands that secrete mucin" [1]. "Hyperplasia and hamartoma of Brunner's glands are benign proliferative lesions of the duodenum. Hyperplasia of Brunner's glands is typically a lesion less than 0.5 cm, characterized by neutral mucin-secreting glands occupying at least 50.0% of the duodenal mucosa in a biopsy specimen" [2,3]. In contrast, hamartoma of Brunner's glands, also known as Brunner gland adenoma, is generally a lesion larger than 0.5 cm [4,5], involving proliferative glands within the duodenal submucosa, mixed with dilated cystic glands and smooth muscle proliferation [6]. The fundamental difference lies in the mixture of other benign components (smooth muscle fibers) with the glands to qualify as an "hamartoma".

"The majority of Brunner gland hamartomas are isolated pedunculated polyps, with a minority being sessile. Their diameter ranges from 1.0 to 2.0 cm, rarely exceeding 5.0 cm, and up to 12.0 cm in some cases" [7,8]. "Most lesions are located in the proximal duodenum, with occurrence decreasing with distance from the pyloric ring: 57.0% in the duodenal bulb, 27.0% in the descending part, and 7.0% in the horizontal part" [9].

"Brunner gland hamartoma is an extremely rare tumor, with an estimated incidence of 0.008% in a single series of 215,000 autopsies" [1].


In this article, we report a clinical case of a patient with a Brunner gland hamartoma of the duodenum who presented with intestinal obstruction and was treated with surgical resection of the polyp via duodenotomy.

## 2. CASE PRESENTATION

He is a 22-year-old young man with no notable medical history. He was admitted with high intestinal obstruction without signs of gastrointestinal bleeding.

Clinical examination and laboratory tests were normal. An abdominal computed tomography (CT) scan showed a tissue lesion in the bulboduodenal region measuring 57mm x 34mm x 22mm, which enhanced moderately after injection of iodinated contrast.


The patient was taken to the operating room after preparation. During exploratory surgery, a duodenal polyp was identified (Fig. 1). The patient underwent surgical resection of the polyp via duodenotomy (Figs. 2 and 3). Postoperative recovery was uneventful.



**Fig. 1. Operative view of duodenal polyp**



**Fig. 2. View after resection of duodenal polyp**



**Fig. 3. Surgical specimen**

Histopathological examination of the surgical specimen revealed a hamartoma of the Brunner's glands, measuring 4.5 cm.

During a 6-month follow-up, the patient remained asymptomatic. A follow-up esophagogastroduodenoscopy showed no residual lesions.

### 3. DISCUSSION

"Brunner's gland hamartoma tends to be predominant in the fifth or sixth decade of life

with an equal distribution between sexes" [7]. The clinical presentation varies; however, the majority of cases are asymptomatic or manifest with nonspecific and vague symptoms such as abdominal pain, nausea, or bloating.

"In symptomatic patients, the most common clinical manifestations are gastrointestinal bleeding (37%) and obstructive symptoms (37%)" [7]. "Gastrointestinal bleeding most often presents as chronic blood loss leading to iron deficiency and anemia" [5]. "Less frequently, when erosions or ulcerations of the tumor occur,

patients may present with melena or hematemesis. These observations are typically described in Brunner's gland hamartomas located beyond the first portion of the duodenum, likely because these lesions are subject to more stress and vascular damage due to gastrointestinal motility" [5]. In our case, the patient was admitted for intestinal obstruction.

"The etiology and pathogenesis of Brunner's gland hamartoma remain unclear. Due to the "anti-acid" function of Brunner's glands, increased acid secretion has been postulated to stimulate these structures to undergo hyperplasia" [3]. Franzin et al. [6] reported an association between Brunner's gland hamartoma and hyperchlorhydria in patients with chronic gastric erosions and duodenal ulcers, but Spellberg et al. [8] did not observe regression of the lesion with acid secretion inhibitors [8]. A second hypothesis suggests an inflammatory origin due to dense infiltration of inflammatory cells [10]. Since lymphocytes are usually present in the normal submucosa of the intestinal tract, the presence of inflammatory foci in Brunner's gland hamartoma is not sufficient to support the inflammatory hypothesis. Finally, it has been suggested that *H. pylori* infection could play a role in the pathogenesis of Brunner's gland hamartoma. In a recent study of 19,100 subjects, *H. pylori* infection was found in five out of seven cases of Brunner's gland hamartoma (71%) [11]. In our patient, no *H. pylori* infection was found. The extremely rare occurrence of Brunner's gland hamartoma and the high prevalence of *H. pylori* infection in the general population do not allow us to establish a clear pathogenic link.

"The duodenal bulb is the most frequent location of Brunner's gland hamartoma (57%)" [12]. "In most cases, these lesions develop into a polypoid mass, usually pedunculated (88%), measuring between 1 and 2 cm" [13].

"Diagnosis is not always straightforward. Radiological findings are often non specific [14]. Indeed, duodenal filling defects can mimic several other lesions such as leiomyoma, lipoma, lymphoma, aberrant pancreatic tissue, or carcinoid tumors" [15]. "CT scan is only useful to confirm the absence of extraluminal extension of Brunner's gland hamartoma" [15].

"Diagnosis can be confirmed by histological examination of the excised mass. Traditional endoscopy with pinch biopsies is generally

negative because pinch biopsies cannot reach tumor tissue located entirely in the submucosal layer" [16].

"Endoscopic or surgical removal of Brunner's gland hamartoma has been suggested to prevent the development of complications (bleeding, severe anemia, intestinal obstruction, or intussusception). Endoscopic polypectomy is the preferred approach, being more cost-effective and less invasive than abdominal surgery" [17,18]. "However, success depends on the location and size of the hamartoma as well as the presence of a pedicle. Several cases of successful endoscopic resections have been reported" [14,19-20]. In our case, the patient had a 57 mm duodenal process complicated by intestinal obstruction, so surgical treatment was necessary [21,22]. Duodenotomy resection of the duodenal polyp was performed without complication, and there was no recurrence of the lesion during the 6-month follow-up [23].

#### 4. CONCLUSION

The Brunner's gland hamartoma is a rare benign tumor of the duodenum with nonspecific clinical manifestations. Its mechanisms remain unclear. Malignant transformation should be cautiously evaluated through histology combined with immunohistochemistry, especially if the lesion size increases and its morphology changes. An observation strategy is employed in the majority of Brunner's gland hamartoma cases. If necessary or in cases of complications, endoscopic and/or surgical resection is required.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

#### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Botsford TW, Crowe P, Croker DW. Tumors of the small intestine. A review of experience with 115 cases including a report of a rare case of malignant hemangio-endothelioma. *Am J Surg.* 1962; 103: 358-365
2. Dhouha B, Ahlem L, Sana BS, et al. Unexpected cause for duodenal obstruction: Brunner's gland hyperplasia. *Pathologica.* 2017;109(4):414-7.
3. Chong KC, Cheah WK, Lenzi JE, et al. Benign duodenal tumors. *Hepatogastroenterology.* 2000;47(35): 1298-300.
4. Mayoral W, Salcedo JA, Montgomery E, et al. Biliary obstruction and pancreatitis caused by Brunner's gland hyperplasia of the ampulla of Vater: a case report and review of the literature. *Endoscopy.* 2000; 32(12):998-1001.
5. Patel ND, Levy AD, Mehrotra AK, et al. Brunner's gland hyperplasia and hamartoma: Imaging features with clinicopathologic correlation. *AJR Am J Roentgenol.* 2006;187(3):715-22.
6. Brosens LAA, Jansen M, Giardiello FM, et al. Polyps of the small intestine. *Diagn Histopathol.* 2011;17(2):69-79.
7. Rocco A, Borriello P, Compare D, et al. Large Brunner's gland adenoma: Case report and literature review. *World J Gastroenterol.* 2006;12(12):1966-8.
8. Gao YP, Zhu JS, Zheng WJ. Brunner's gland adenoma of duodenum: A case report and literature review. *World J Gastroenterol.* 2004;10(17):2616-7.
9. Walden DT, Marcon NE. Endoscopic injection and polypectomy for bleeding Brunner's gland hamartoma: Case report and expanded literature review. *Gastrointest Endosc.* 1998;47(5):403-7.
10. Kim K, Jang SJ, Song HJ, et al. Clinicopathologic characteristics and mucin expression in Brunner's gland proliferating lesions. *Dig Dis Sci.* 2013;58(1):194-201.
11. Krishnamurthy P, Junaid O, Moezzi J, et al. Gastric outlet obstruction caused by Brunner's gland hyperplasia: Case report and review of literature. *Gastrointest Endosc.* 2006;64(3):464-7.
12. Bakir MA, AlYousef MY, Alsohaibani FI, et al. Brunner's glands hamartoma with pylorusobstruction: A case report and review of literature. *J Surg Case Rep.* 2020;8:rjaa191.
13. Botsford TW, Crowe P, Crocker DW. Tumors of the small intestine. A review of experience with 115 cases including a report of a rare case of malignant hemangio-endothelioma. *Am J Surg.* 1962; 103: 358-65.
14. Iusco D, Roncoroni L, Violi V, et al. Brunner's gland hamartoma: "overtreatment" of a voluminous mass simulating a malignancy of the pancreaticoduodenal area. *Jop.* 2005;6(4): 348-53.
15. Spellberg MA, Vucelic B. A case of Brunner's glands hyperplasia with diarrhea responsive to cimetidine. *Am J Gastroenterol.* 1980;73(6):519-22.
16. Destek S, Gul VO. Brunner's gland hyperplasias and hamartomas in association with *Helicobacter pylori*. *Can J Gastroenterol Hepatol.* 2019;2019 (6340565):6.
17. Sakurai T, Sakashita H, Honjo G, et al. Gastric foveolar metaplasia with dysplastic changes in Brunner gland hyperplasia: possible precursor lesions for Brunner gland adenocarcinoma. *Am J Surg Pathol.* 2005;29(11): 1442-8.
18. Kovacevic I, Ljubivic N, Cupic H, et al. *Helicobacter pylori* infection in patients with Brunner's gland adenoma. *Acta Med Croatica.* 2001;55(4-5):157-60.
19. Stolte M, Schwabe H, Prestele H. Relationship between diseases of the pancreas and hyperplasia of Brunner's glands. *Virchows Arch A Pathol Anat Histol.* 1981;394(1-2):75-87.
20. Akaki M, Taniguchi S, Hatakeyama K, et al. Duodenal mucosal damage is associated with proliferative activity of Brunner's gland hamartoma: A case report. *BMC Gastroenterol.* 2014;14: 14.
21. Mumtaz R, Shah IA, Ramirez FC. "Brunner's gland hamartoma simulating a pancreatic mass with duodenal obstruction. *Gastrointest Endosc.* 2002; 56:932-4
22. Delvecchio K, Seman S. Successful surgical excision of a massive

- symptomatic partially obstructing  
Brunner's gland hamartoma: A case  
report. J Surg Case Rep. 2016;2016  
(12)
23. Dhali A, Ray S, Mandal TS, et al. Surgical  
Outcome of Brunner's Gland Hamartoma:  
A Single-Centre Experience. Surg J (N Y).  
2022;8(1)

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/123210>