



# Characterization of Extended Spectrum Beta-lactamase Uropathogens Isolated from Refugees with Urinary Tract Infections in Nakivale Refugee Settlement Camp, Southwestern Uganda

Ayan Ahmed Hussein <sup>a</sup>, Kennedy Kassaza <sup>a</sup>,  
James Mwesigye <sup>a</sup>, Nalumaga Pauline Petra <sup>a</sup>,  
Bashir Mwambi <sup>b</sup>, Taseera Kabanda <sup>a</sup> and Joel Bazira <sup>a\*</sup>

<sup>a</sup> Department of Microbiology, Faculty of Medicine, Mbarara University of Science and Technology, P.O Box 1410, Mbarara, Mbarara city, Uganda.

<sup>b</sup> Department of Medical Laboratory Science, Institute of Allied Health Sciences, Clarke, International University P.O Box 7782 Kampala, Uganda.

## Authors' contributions

This work was carried out in collaboration among all authors. Author AAH conceptualized the study and participated in data collection. Authors KK and JM contributed to the design and conduct of the molecular assays. Author JM carried out the culture and sensitivity. Author TK contributed to study design and analysis. Author BM participated in data analysis. Author JB over saw the overall running of the study. Authors NPP and BM wrote the final manuscript. All authors read and approved the final version of the manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ISRR/2024/v13i2181>

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125158>

**Original Research Article**

**Received: 20/08/2024**  
**Accepted: 22/10/2024**  
**Published: 05/11/2024**

\*Corresponding author: Email: [jbazira@must.ac.ug](mailto:jbazira@must.ac.ug);

**Cite as:** Hussein, Ayan Ahmed, Kennedy Kassaza, James Mwesigye, Nalumaga Pauline Petra, Bashir Mwambi, Taseera Kabanda, and Joel Bazira. 2024. "Characterization of Extended Spectrum Beta-Lactamase Uropathogens Isolated from Refugees With Urinary Tract Infections in Nakivale Refugee Settlement Camp, Southwestern Uganda". *International STD Research & Reviews* 13 (2):67-78. <https://doi.org/10.9734/ISRR/2024/v13i2181>.

## ABSTRACT

The World Health Organization estimates one in four individuals has had at least one urinary tract infection (UTI) episode requiring treatment with an antimicrobial agent. At Nakivale refugee camp, the overwhelming number of refugees often associated with poor living conditions predispose the refugees to urinary tract infections.

This study determined the prevalence of UTIs, the antimicrobial susceptibility pattern of the isolated bacterial pathogens, the prevalence of Extended-Spectrum Beta-Lactamase (ESBL) bacteria, and the molecular characterization of genes encoding ESBLs among refugees in the Nakivale refugee settlement.

This was a cross-sectional study that involved 216 outpatients who visited Nakivale Health Centre III between July and September 2020. The urine samples were received and examined at the microbiology laboratory of Mbarara University of Science and Technology. The urine samples were cultured and identified. Antibiotic susceptibility was carried out following CLSI recommended guidelines while the presence of genes encoding ESBL was detected using conventional PCR amplification.

The prevalence of UTI was 24.1% (52/216). *Staphylococcus aureus* was the most prevalent causative agent, accounting for 22/52 (42.31%) of total isolates, followed by *Escherichia coli* 21/52(40.38%). Multidrug-resistant isolates accounted for 71.15% (37/52). A total of twenty-one isolates (70.0%) were extended spectrum beta-lactamase producing bacteria. The most prevalent genes were TEM beta-lactamase (blaTEM) and CTX-M beta-lactamase (blaCTX-M).

The prevalence of UTI among refugees in the Nakivale settlement was high which calls for continuous epidemiological surveys to determine the prevalence of multi-drug resistance uropathogens including ESBL-producing organisms across refugee camps in Uganda.

**Keywords:** Uropathogens; extended beta-lactamase organisms; nakivale refugee settlement camp.

## 1. INTRODUCTION

Urinary Tract Infections are infections of the urinary tract system that arise from bacterial migration within the genitourinary tract. They are usually transmitted through contaminated hands, normal flora migration, or through sexual intercourse [1]. Globally, the incidences of UTI have become endemic which has significantly increased the use of antimicrobial agents [2]. The widespread use of these antibiotics has caused the emergence and spread of resistant bacteria including extended-spectrum beta-lactamase-producing organisms [3]. Extended-spectrum beta-lactamase producing uropathogens are widely spread around the world, and their severity is higher in poorer nations like Uganda [4]. ESBL-producing organisms frequently exhibit characteristics of multidrug resistance [5] to frequently used antibiotics because they frequently carry genes *blaTEM*, *blaCTX-M*, and *blaSHV* encoding resistance to aminoglycosides thus, ESBL-producing organisms often possess multidrug resistance phenotypes [5].

Uganda is Africa's largest refugee-hosting country and the 8<sup>th</sup> in the world with a total

population of about 1,381,122 of which 130,462 (9.45%) are found in the Nakivale refugee camp [6]. The overwhelming number of refugees is often associated with poor living conditions such as communal bathrooms and toilets and multiple sex partners resulting in compromised health [7]. As a result, a high number of suspected urinary tract infections are reported at the Nakivale refugee camp (at least 20 suspected UTIs per day) according to the record from the health centre.

According to unpublished Health Center III records in the Nakivale refugee settlement, the prevalence of clinically diagnosed UTI was seen to be 15%. Resistant UTIs are among the challenges faced by healthcare providers in the refugee population [8]. Accordingly, with the influx of refugees, new urinary tract infections caused by gram-negative bacterial-resistant strains could be introduced as already reported by some studies [9]. Delayed recognition of severe UTI caused by ESBL-producing organisms and inappropriate treatment with antibiotics has been associated with increased mortality and morbidity in resettlement camps a phenomenon that stresses the need for evaluation of UTI etiology.

## 2. MATERIALS AND METHODS

### 2.1 Study Population and Sample Collection

The study was a descriptive cross-sectional study, conducted among 216 outpatients attending Nakivale Health Centre III in Nakivale Health Centre III refugee settlement in Isingiro district located 59km away from Mbarara district, South-western from July 2020 to September 2020. Participants with complaints suggestive of UTI, such as dysuria, frequency, urgency, suprapubic pain, and cloudy urine as identified by the attending clinician, were considered for the study after receiving informed consent. Participants who did not provide a urine sample and females who were in their menstrual cycle were excluded. Participants were given oral and written instructions to collect midstream urine in sterile leak-proof labeled urine containers.

### 2.2 Laboratory Tests, Urine Cultures, and Isolation of Organisms

The urine samples were collected from the Nakivale settlement camp transported in a cold chain and received and examined at the microbiology laboratory of Mbarara University of Science and Technology. The color and turbidity of the urine samples were recorded. The urine was microscopically wet-prepared to look for significant pyuria (>10 cells/ $\mu$ l), casts, red cells, bacteria, and yeast. Urine dipstick biochemical testing and primary gram staining were performed. The urine samples were inoculated using a 2mm standard loop and plated on Cystine Lactose Electrolyte-Deficient (CLED) agar with bromothymol blue (HiMedia Laboratories supplies) and incubated aerobically at 37°C overnight for 24 hours. Colonies were counted 50 and above colony forming units which were equivalent to 10000 ( $10^4$ ) by ml (CFU/mL) were significant [10]. Purity plating was done on MacConkey Agar, Blood Agar, and Mannitol Salt Agar (Oxoid).

The pure isolated bacterial colonies were identified using gram staining, colony morphology [11], Biochemical testing that included citrate utilization test, urease test, triple sugar iron agar (TSI) test, indole test, sulfur indole motility [12], and Analytical Profile Index (API).

### 2.3 Antimicrobial Susceptibility Testing

The antimicrobial susceptibility pattern of the isolated bacterial pathogens was performed using the disc diffusion method on Mueller-Hinton agar according to the guidelines of the Clinical and Laboratory Standards Institute [13]. Antibiotic discs from Oxoid that were used for both gram-positive and gram-negative bacteria included: Nitrofurantoin (100  $\mu$ g), Ampicillin (10 $\mu$ g), Ciprofloxacin (5 $\mu$ g), Ceftriaxone (30 $\mu$ g), Amoxicillin-clavulanic acid (20/10 $\mu$ g), and Chloramphenicol (30 $\mu$ g) while Erythromycin (15 $\mu$ g), and Penicillin (10 $\mu$ g) were used for only gram-positive bacteria.

### 2.4 Screening for ESBLs

Gram-negative bacteria were confirmed for the presence of extended-spectrum  $\beta$ -lactamases by using the phenotypic detection method as per CLSI guidelines. The discs used were Ceftazidime (30 $\mu$ g) and ceftazidime-clavulanic acid (30/10 $\mu$ g), cefotaxime (30 $\mu$ g) and cefotaxime-clavulanic acid (30/10 $\mu$ g). The isolates were interpreted as ESBL producers if a clear extension of the edge of the inhibition zone of cephalosporin towards the disc of cephalosporin with clavulanic acid was observed [14]. Reference strains of *Escherichia coli* ATCC25922 (ESBL-negative), *Klebsiella pneumoniae* ATCC 700603 (ESBL-positive), and *Staphylococcus aureus* ATCC 29213. *Pseudomonas aeruginosa* ATCC 27853, *Enterococcus faecalis* ATCC 29212 were used [15].

### 2.5 Genotypic Characterization of ESBLs

ESBL phenotypes were sub-cultured at 37°C overnight in Muller Hinton Agar (Oxoid, Wade Road, Basingstoke, UK). The bacterial DNA was extracted using the Zymo-Research quick-g-DNA Miniprep kit and protocol, v1.0.0.

The presence of genes encoding ESBL was detected using PCR amplification. This study detected *bla*-SHV, *bla*-CTX-M [16], and *bla*-TEM [17], genes. The PCR master mix reagents for each gene target were as follows: 12.5  $\mu$ L master mix consisting of One Taq quick-load 2x master mix/w standard buffer, dNTPs & Taq polymerase (M0486S), 1.5  $\mu$ L forward (100  $\mu$ M), 1.5  $\mu$ L primary reverse (100  $\mu$ M), 5  $\mu$ L DNA template and RNAase-free water up to 25  $\mu$ L.

**Table 1. Primers sets used in the detection of ESBL encoding genes from the study isolates [18]**

No.	Resistance gene	Primer	Annealing temperature	Amplicon bp
1	<i>bla</i> TEM	(F) 5'-ATTTCCGTGTCGCCCTTAT-3 (R) 5'-CTACGATACGGGAGGGCTTA-3'	54.5°C	759
2	<i>bla</i> CTX-M	(F) 5'-ATGATGAAAAAATCGTTATGC-3' (R) 5'-CAGCATCTCCCAGCCTAAT-3'	57°C	489
3	<i>bla</i> SHV	(F) 5'- CCTGTTAGCCACCCTGCC-3' (R) 5'- CCGCAGATAAATCACCCAC-3'	60°C	768

**PCR Cycling:** The PCR process was carried out in a thermocycler (Multigene Optimax) with a denaturation cycle of 95°C for 15 min, DNA amplification of 30 cycles (94°C for 1 min, 54.5-60°C for 1 min, and 72°C for 1 min) and final extension cycle of 72°C for 5 min (See Table 1). DNA Amplicon was electrophoresed using 1.5% agarose gel, in Tris-Borate EDTA buffer (TBE) 1xconcentration, Safe View Classic™ DNA stain (cat # G108), 6x loading dye (Thermo Scientific #R0611), and DNA ladder/marker 100 bp (NEB-Biolabs #N3231L). DNA Bands were visualized on a Dark reader Transilluminator (DR46-B).

## 2.6 Data Management and Analysis

The obtained data were entered in Microsoft Excel, and imported to STATA version 12.0 for analysis. Evaluations were carried out at a 95 %

confidence level (95% CI). Results were presented in the form of tables, and pie charts.

## 3. RESULTS

### 3.1 The Demographic Characteristics of Study Participants

There were 216 participants in the study, including 164 (75.93%) females and 52 (24.07%) males. Participant ages ranged from 18 to 76, with an average age of 35.86 (SD=11.218) and a median of 34. 86 (39.81%). Democratic Republic of the Congo received the greatest number of refugees 86(39.81%) followed by those from Somalia 58 (26.85%). Regarding education, 89 (41.20%) of the participants were not educated, followed by those whose level of education was primary accounting for 82(37.96%) as shown in Table 2.

**Table 2. Demographic characteristics of the study participants**

Variable	Frequency (n)	Percentage
<b>Sex</b>		
Female	164	75.93 (69.66-81.47)
Males	52	24.07 (18.54 - 30.34)
<b>Age group</b>		
18-76 years	216	100 (98.31 - 100)
<b>Country of origin</b>		
Burundi	27	12.50 (8.40 - 17.66)
Democratic Republic of Congo	84	38.89 (32.35 - 45.74)
Ethiopia	01	0.46 (0.012 - 2.55)
Rwanda	35	16.20 (11.55 - 21.80)
South Sudan	11	5.09 (2.57 - 8.93)
Somalia	58	26.85 (21.07 - 33.29)
<b>Education level</b>		
None	89	41.20 (34.57 - 48.08)
Primary	82	37.96 (31.47 - 44.79)
Secondary	37	17.13 (12.37 - 22.83)
Tertiary	8	3.70 (1.61 - 7.17)
<b>Marital status</b>		
Divorced	15	6.94 (3.94 - 11.19)
Married	163	75.46 (69.17 - 81.04)
Single	27	12.50 (8.40 - 17.66)
Widowed	11	5.09 (2.57 - 8.93)

### 3.2 The Prevalence of Urinary Tract Infections among Refugees in the Nakivale Refugee Settlement

The overall prevalence of UTI was 24.1% (52/216). The prevalence of UTI was 25% (41/52) among females and 21.15% (11/52) among males. With country-of-origin South Sudan had the highest prevalence of UTI which was 36.36% (4/11), followed by the Democratic Republic of the Congo at 29.76% (25/84). Refugees of primary level of education had the highest prevalence of UTI 29.27% (24/84), followed by uneducated 23.60% (21/89). Lastly, single refugees had the highest prevalence of UTI 29.63% (8/27), followed by married refugees 23.93% (39/163) (Table 3).

### 3.3 The Bacterial Etiology of Urinary Tract Infections among Refugees in Nakivale Settlement

*Staphylococcus aureus* was the most common organism accounting for 22/52 (42%) of total isolates, followed by *Escherichia coli* 21/52

(40%), *Klebsiella pneumoniae* 4/52 (8%), *Salmonella arizonae* 2/52 (4%) while *Citrobacter freundii*, *Hafnia species*, and *Pseudomonas aeruginosa* accounted to 1/52 (2%) each as illustrated in Fig. 1.

### 3.4 The Antimicrobial Susceptibility Pattern of the Isolated Bacterial Pathogens among Refugees in the Nakivale Settlement

The most common bacterial isolates that are *Staphylococcus aureus* were most sensitive to Nitrofurantoin (86.36%) and Amoxicillin/Clavulanic Acid (81.82%), Chloramphenicol (72.73%), and Ciprofloxacin (63.64%) but most resistant to Penicillin (95.45%), Ceftriaxone (68.18%) and Erythromycin (63.64%). While *Escherichia coli* was most sensitive to Chloramphenicol (90.48%) and Nitrofurantoin (76.19%) but most resistant to Amoxicillin/Clavulanic Acid (90.48%), Ampicillin (90.48%), Ceftriaxone (76.19%), and Ciprofloxacin (61.90%). See Table 4.

Table 3. The prevalence and demographic distribution of UTI

Sex	Negative cultures for bacteria (%)	Positive cultures for bacteria (%)	Total	Prevalence of UTI % (95% CI)
<b>Total enrolment</b>	<b>164</b>	<b>52</b>	<b>216</b>	<b>24.07 (18.53 - 30.34)</b>
<b>Positivity by Gender</b>				
Female	123	41	164	25.00 (18.58 - 32.35)
Male	41	11	52	21.15 (11.06 - 34.70)
<b>Country of origin</b>				
Burundi	20	7	27	25.93 (11.11 - 46.28)
Democratic Republic of the Congo	59	25	84	29.76 (20.27 - 40.73)
Rwanda	30	5	35	22.7 (4.81 - 30.26)
South Sudan	7	4	11	36.36 (10.93 - 69.21)
Somalia	47	11	58	18.96 (9.87 - 31.41)
<b>Education level</b>				
None	68	21	89	23.60 (15.24 - 33.78)
Primary	58	24	82	29.27 (19.73 - 40.35)
Secondary	31	6	37	16.22 (6.19 - 32.01)
Tertiary	7	1	8	12.50 (0.32 - 52.65)
<b>Marital status</b>				
Divorced	12	3	15	20.00 (16.58 - 40.46)
Married	124	39	163	23.93 (17.60 - 31.22)
Single	19	8	27	29.63 (13.75 - 50.18)
Widowed	9	2	11	18.18 (2.28 - 51.77)




Fig. 1. The bacterial etiology of UTI among refugees in Nakivale settlement

### 3.5 Phenotypic Characterization of Extended Spectrum Beta-Lactamase Organisms among Refugees in the Nakivale Refugee Settlement

Phenotypic characterization of ESBL production was performed on 30 Gram-negative organisms and 21 (70.00%) were ESBL producers for the different organisms while 9(30%) were non-ESBL producers. (see Table 5).

### 3.6 Genotypic Characterization of Extended Spectrum Beta-Lactamase Organisms among Refugees in Nakivale Refuge

The presence of any genes of interest was detected in 8 (26.67%) of 30 Extended Spectrum Beta-Lactamase organisms. The most prevalent genes were *bla*TEM and *bla*CTX-M, both of which were found in 5 (62.5%) of the organisms possessing genes while *bla*SHV was found in three (37.5%) of the organisms containing genes, as indicated in Table 6.

## 4. DISCUSSION

### 4.1 The Prevalence of Urinary Tract Infections

The prevalence of UTI was 24.1% in this study, this finding was consistent with the findings of Tibyangye [19] study in Bushenyi, where the prevalence of UTI was (22.5%) and Odongo [20] study in Gulu, where the prevalence was (24.2%). The rationale for the consistent results

could be due to geographic locations as indicated by Tandogdu, [21].

When compared to Turkman's study of refugees [22], Aro's study of refugees in Finland [23], Gulati's study of Syrian refugees in Asia [24], and UNHCR's study of refugees in Tanzania [25], the finding was high according to the Uganda refugee health report 2018, UTI constituted 2% of all consultations made in all refugee settlements, a figure contradicts the findings of this study. Thus, the cause for disparity in UTI prevalence may be attributed to the procedures employed to diagnose UTI. We employed urine culture in our study, whereas they used conventional urinalysis in other studies.

However, the study prevalence was low when compared to 38.8% in a study conducted by Kabugo [26] at Mulago Hospital in Uganda and 32.2% in a study conducted among patients attending hospitals in Bushenyi [27]. Although our frequency is not much lower than that of Mulago Hospital and KIU Teaching Hospital, this is because our study recruited largely outpatients who came to lesser health units rather than patients who sought care at a tertiary hospital.

The high prevalence of UTI could be attributed to the large number of refugees who are frequently connected with poor living conditions such as communal restrooms and toilets as well as several sex partners, resulting in compromised health [28].

**Table 4. Antibiotic susceptibility profile for individual antibiotics**

Bacteria Isolate	Susceptibility status	Antimicrobial Susceptibility results (%)								Total Isolated
		CIP	AUG	CXN	CAF	NIT	AMP	E	P	
<b>Gram Positive</b>										
<i>S.aureus</i>	Resistant	8 (36.36)	4 (18.18)	15 (68.18)	6 (27.27)	3 (13.64)	13 (59.09)	14 (63.64)	21 (95.45)	22
	Sensitive	14 (63.64)	18 (81.82)	7 (31.82)	16 (72.73)	19 (86.36)	9 (40.91)	8 (36.36)	1 (4.55)	
<b>Gram-Negative</b>										
<i>E. Coli</i>	Resistant	13 (61.90)	19 (90.48)	16 (76.19)	2 (9.52)	5 (23.81)	19 (90.48)	-	-	21
	Sensitive	8 (38.10)	2(9.52)	5(23.81)	19 (90.48)	16 (76.19)	2(9.52)	-	-	
<i>Klebsiella pneumoniae</i>	Resistant	0(0.00)	4(100)	2(50)	3 (75.000)	4(100)	4(100)	-	-	4
	Sensitive	4(100)	0(0.00)	2(50)	1 (25.00)	0(0.00)	0(0.00)	-	-	
<i>Salmonella arizonae</i>	Resistant	2(100)	2(100)	2(100)	0(0.00)	0(0.00)	2(100)	-	-	2
	Sensitive	0(0.00)	0(0.00)	0(0.00)	2(100)	2(100)	0(0.00)	-	-	
<i>Citrobacter freundii</i>	Resistant	0(0.00)	1 (100)	1 (100)	0(0.00)	0(0.00)	1(100)	-	-	1
	Sensitive	1 (100)	0(0.00)	0(0.00)	1(100)	1(100)	0(0.00)	-	-	
<i>Hafnia spp</i>	Resistant	1 (100)	1(100)	1(100)	0(0.00)	1(100)	1(100)	-	-	1
	Sensitive	0(0.00)	0(0.00)	0(0.00)	1(100)	0(0.00)	0(0.00)	-	-	
<i>Pseudomonas aeruginosa</i>	Resistant	1(100)	1(100)	0(0.00)	1(100)	1(100)	1(100)	-	-	1
	Sensitive	0(0.00)	0(0.00)	1(100)	0(0.00)	0(0.00)	0(0.00)	-	-	
<b>Total</b>	Resistant	25 (48.08)	32 (61.54)	37 (71.15)	12 (23.08)	14 (26.92)	41 (78.85)	-	-	52
	Sensitive	27 (51.92)	20 (38.46)	15 (28.85)	40 (76.92)	38 (73.08)	11 (21.15)	-	-	

CIP= Ciprofloxacin; AUG=Amoxicillin/ Clavulanic Acid; CXN=Ceftriaxone; CAF=Chloramphenicol; NIT=Nitofurantoïn; AMP=Ampicillin; E=Erythromycin; P=Penicillin; -=Not tested

**Table 5. ESBL production by the different organisms**

Organisms isolated	ESBL producer		Total
	No (%)	Yes (%)	
<i>Escherichia coli</i>	7(33.33)	14(66.67)	21
<i>Klebsiella pneumoniae</i>	1(25.00)	3(75.00)	4
<i>Salmonella arizonae</i>	1(50.00)	1(50.00)	2
<i>Citrobacter freundii</i>	n.d	1(100%)	1
<i>Hafnia species</i>	n.d	1(100)	1
<i>Pseudomonas aeruginosa</i>	n.d	1(100)	1
<b>Total</b>	<b>9(30.00)</b>	<b>21(70.00)</b>	<b>30</b>

**Table 6. Genotypic characterization of extended spectrum beta-lactamase organisms**

Organisms isolated	ESBL producer			Positive for any gene (%)	Negative for any gene (%)	Total (%)
	Genes Isolated					
	<i>bla</i> TEM	<i>bla</i> SHV	<i>bla</i> CTX-M			
<i>Escherichia coli</i>	2	2	4	5 (23.81)	16(76.19)	21(100)
<i>Klebsiella pneumoniae</i>	2	1	0	2(50.00)	2(50.00)	4(100)
<i>Pseudomonas aeruginosa</i>	1	0	1	1(100.00)	n.d	1(100)
<b>Total</b>	<b>5</b>	<b>3</b>	<b>5</b>	<b>8(26.67)</b>	<b>22(73.33)</b>	<b>30 (100)</b>

Although numerous studies have found *Escherichia coli* was the most prevalent bacteria causing UTIs [29,30], the most common organism in our study was *Staphylococcus aureus*. *Staphylococcus aureus* was the most prevalent bacterial isolate in this investigation, accounting for 42.31% of total isolates, followed by *Escherichia coli* (40.38%). This was consistent with Ekwealor's [31] study in Nigeria, which discovered that the most prevalent isolate was *Staphylococcus aureus* (28%), followed by *Escherichia coli* (24.6%). Research conducted in Ethiopia by Regea [32] discovered that the most prevalent isolates were, *Staphylococcus aureus* (24.2%), *CoN Staphylococcus spp* (24.2%), *E.coli* (12.1%), as well as *K. pneumoniae* (12.1%).

#### 4.2 Antimicrobial Susceptibility Pattern of the Individual Isolated Bacterial Pathogens among Refugees in the Nakivale Settlement

The sensitivity of *Staphylococcus aureus* in this study was in line with a study done by Bitew et al., [33] where almost all gram-positive sensitive to nitrofurantoin (97.1%) and a study done by Anejo-Okopi and colleagues in 2017 [34] who also found *Staphylococcus aureus* to be sensitive to Ciprofloxacin (83.33%),

Chloramphenicol (75%) and Amoxicillin (83.33%).

Regarding resistance of *Staphylococcus aureus*, this study was consistent with a study done by Bitew et al. [33] who found that gram-positive was resistant to erythromycin (82.2%), a study by Gebremariam et al. [22] which found that Gram-positive bacterial isolates were highly resistant to ampicillin (81.5%).

Furthermore, the study findings of the sensitivity of *E. coli* differ from results from a study done in Nigeria by Anejo-Okopi et al. [34] which showed that *Escherichia coli* was more sensitive to Chloramphenicol and Nitrofurantoin but instead more sensitive to Ciprofloxacin (60%), Streptomycin (100%), and 70% sensitive to chloramphenicol. A study done by Mwaka et al. [23] at Mulago Hospital in Uganda also found that bacteria isolated were sensitive to nitrofurantoin (98.3%). This shows that nitrofurantoin can still be used as a first-line antibiotic in UTIs.

The prevalence of ESBL in this study was high when compared with the overall prevalence of ESBL in Kenya (37.4%), Rwanda (38.3), Algeria (19.9%) (29), and 6.4% in the United States [25]. Another study by Mansouri in 2019 [35] determined increased ESBL production by



69.1–78.6% in Europe. Increased prevalence (55.5%) of ESBL-positive organisms has also been noted in Nairobi Kenya by Magale in 2015 [36] and between 27.8% to 64.9% in Uganda [29-31].

Therefore, this high prevalence of ESBL-producing *Enterobacteriaceae* reflects a more serious public health problem. Dissemination of ESBLs compromises the activity of broad-spectrum antibiotics creating major therapeutic difficulties with a significant impact on the outcomes for patients [32]. Additionally, a study by Fatima [37] in Pakistan reported that 97% of ESBL-producing bacteria isolates were resistant to Ampicillin, therefore, Ampicillin resistance in this study could have resulted from the high prevalence of ESBL-producing *Enterobacteriaceae*. Furthermore, the high prevalence of ESBL-producing bacteria isolates could also be the reason for the high multi-drug resistance in this study.

In this study, the most common genes were *bla*TEM and *bla*CTX-M each detected in 5 (62.5%) of the organisms with genes while *bla*SHV was detected in 3 (37.5%) of the organisms with genes. This contradicts the finding in studies done by Jena [38,39] where in all these studies *bla*TEM the only predominant gene was followed by *bla*CTX-M. Other contradictions were reflected by studies conducted by Seyedjavadi [40] and Ahmed [41] where in their case *bla*CTX-M was the predominant gene followed by *bla*TEM. The reason for such could be the difference in study areas and people.

Despite the difference on which gene is common the percentage of *bla*TEM and *bla*CTX-M in this study and others was above 60%. Thus, it can be conclusive that the most common genes are *bla*TEM and *bla*CTX-M. However, Sequencing of *bla*TEM and *bla*CTX-M was not performed as well as determining pandemic clones ST131 and ST405 of *E. coli* that would further determine the disease burden [42].

## 5. CONCLUSION

The prevalence of UTI among refugees in Nakivale settlement is high. *Staphylococcus aureus* is the main cause of UTI followed by *Escherichia coli* and *Klebsiella pneumoniae*.

Multidrug resistance was 71.15% in tested isolates, whereas ESBL production in *Enterobacterales* was 70%. The common ESBL genes detected were *bla*TEM and *bla*CTX-M. Conducting continuous epidemiological surveys to determine the existence of ESBL-producing organisms across refugee camps in Uganda.

## DISCLAIMER

This paper is an extended version of a preprint document of the same author.

The preprint document is available in this link: [https://www.researchgate.net/publication/360351544\\_Prevalence\\_and\\_Genotypic\\_Characterization\\_of\\_Extended\\_Spectrum\\_Beta-Lactamase\\_Uropathogens\\_Isolated\\_from\\_Refugees\\_with\\_Urinary\\_Tract\\_Infections\\_in\\_Nakivale\\_Refugee\\_Settlement\\_camp\\_Southwestern\\_Uganda](https://www.researchgate.net/publication/360351544_Prevalence_and_Genotypic_Characterization_of_Extended_Spectrum_Beta-Lactamase_Uropathogens_Isolated_from_Refugees_with_Urinary_Tract_Infections_in_Nakivale_Refugee_Settlement_camp_Southwestern_Uganda)

[As per journal policy, preprint /repository article can be published as a journal article, provided it is not published in any other journal]

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## CONSENT AND ETHICS APPROVAL

This study was approved by the Research and Ethics Committee of Mbarara University of Science and Technology (Ref MUREC 1/7). Upon clearance, permission was sought from the Office of the Prime Minister to research a refugee population, and informed consent was sought from each participant.

## AVAILABILITY OF DATA AND MATERIAL

The datasets used and analyzed during the current study are available from the corresponding author upon reasonable request.

## ACKNOWLEDGEMENT

We appreciate the staff of Nakivale Health Centre III in Nakivale Health Centre III refugee settlement. We are also grateful for the Laboratory staff of Mbarara University of Science and Technology Department of

Microbiology for allowing us to work with them in the study.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Flores-Mireles AL, Walker JN, Caparon M, Hultgren SJJNm. Urinary tract infections: Epidemiology, mechanisms of infection and treatment options. 2015;13(5):269-84.
2. Lee DS, Lee SJ, Choe H-SJBri. Community-acquired urinary tract infection by *Escherichia coli* in the era of antibiotic resistance. 2018;2018.
3. Teklu DS, Negeri AA, Legese MH, Bedada TL, Woldemariam HK, Tullu KDJAR, et al. Extended-spectrum beta-lactamase production and multi-drug resistance among Enterobacteriaceae isolated in Addis Ababa, Ethiopia. 2019;8(1):1-12.
4. Gashaw M, Berhane M, Bekele S, Kibru G, Teshager L, Yilma Y, Ahmed Y, Fentahun N, Assefa H, Wieser A. Emergence of high drug resistant bacterial isolates from patients with health care associated infections at Jimma University medical center: A cross sectional study. *Antimicrobial Resistance & Infection Control*. 2018;7(138).
5. Ventola CL. The antibiotic resistance crisis: Part 1: Causes and threats. *Pharmacy and therapeutics*. P T. 2015;40(4):277-83.
6. Unhcr. UNHCR, Government of Uganda, Office of the Prime Minister, Operational portal refugee situations. Kampala, Uganda; 2019.
7. Gebrehiwet K, Abrha F, Gebreyesus H, Teweldemedhin M. The social health impact of Eritrean refugees on the host communities: The case of May-ayni refugee camp, Northern Ethiopia. *BMC Research Notes*. 2020;13:1-5.
8. Sabih A Lscutiuf. Complicated urinary tract infections. StatPearls Publishing; Treasure Island (FL); 2022.
9. Ursel H, Bernhard K, Karathana M, Niels , Christian Z. Multidrug-resistant bacteria in unaccompanied refugee minors arriving in Frankfurt am Main, Germany, October to November 2015. *Eurosurveillance*. 2016;21(2).
10. Tille P. Bailey & Scott's diagnostic microbiology e-book. Elsevier Health Sciences; 2015.
11. Carter G. Isolation and identification of bacteria from clinical specimens. *Diagnostic procedure in veterinary bacteriology and mycology*: Elsevier. 1990;19-39.
12. Dimri AG, Chaudhary S, Singh D, Chauhan A, Aggarwal M. Morphological and biochemical characterization of food borne Gram-positive and Gram-negative bacteria. *Science Archives*. 2020;1(1):16-23.
13. C. M100 Performance Standards for Antimicrobial Susceptibility Testing; 2018.
14. Andrew B KA, Bazira J. Prevalence of extended-spectrum beta-lactamases-producing microorganisms in patients admitted at KRRH, Southwestern Uganda. Prevalence of extended-spectrum beta-lactamases-producing microorganisms in patients admitted at KRRH, Southwestern Uganda. *Int J Microbiol*; 2017.
15. Musinguzi B, Kabajulizi I, Mpeirwe M, Turugurwa J, Kabanda TJAIID. Incidence and etiology of catheter associated urinary tract infection among admitted patients at Kabale Regional Referral Hospital, South Western Uganda. *Advances in Infectious Diseases*. 2019;9(3).
16. Momtaz H, Rahimi E, Moshkelani SJVM. Molecular detection of antimicrobial resistance genes in *E. coli* isolated from slaughtered commercial chickens in Iran. 2012;57(4):193-7.
17. Wu H, Wang M, Liu Y, Wang X, Wang Y, Lu J, et al. Characterization of antimicrobial resistance in *Klebsiella* species isolated from chicken broilers. 2016;232:95-102.
18. Hayati M, Indrawati A, Mayasari NLPI, Istiyarningsih I, Atikah NJVw. Molecular detection of extended-spectrum  $\beta$ -lactamase-producing *Klebsiella pneumoniae* isolates of chicken origin from East Java, Indonesia. 2019;12(4):578.
19. Tibyangye J, Okech MA, Nyabayo JM, Nakavuma JLJBMJRJ. *In vitro* antibacterial activity of *Ocimum suave* essential oils against uropathogens isolated from patients in selected hospitals in Bushenyi

- district, Uganda. *Br Microbiol Res J*. 2015;8(3):489-98.
20. Odongo CO, Anywar DA, Luryamamoi K, Odongo PJBID. Antibigrams from community-acquired uropathogens in Gulu, northern Uganda-a cross-sectional study. 2013;13:1-8.
  21. Tandogdu Z, Cai T, Koves B, Wagenlehner F, Bjerklund-Johansen TE. Urinary tract infections in immunocompromised patients with diabetes, chronic kidney disease, and kidney transplant. *European urology focus*. 2016;2.
  22. Gebremariam G, Legese H, Woldu Y, Araya T, Hagos K, Gebreyesus Wasihun A. Bacteriological profile, risk factors and antimicrobial susceptibility patterns of symptomatic urinary tract infection among students of Mekelle University, northern Ethiopia. *BMC infectious diseases*. 2019;19:1-11.
  23. Mwaka AD, Mayanja-Kizza H, Kigonya E, Kaddu-Mulindwa D. Bacteriuria among adult non-pregnant women attending Mulago hospital assessment centre in Uganda. *African health sciences*. 2011;11(2).
  24. Storberg V. ESBL-producing Enterobacteriaceae in Africa—a non-systematic literature review of research published 2008–2012. *Infection ecology & epidemiology*. 2014;4(1):20342.
  25. Kaye KS, Gupta V, Mulgirigama A, Joshi AV, Scangarella-Oman NE, Yu K, et al. Antimicrobial resistance trends in urine *Escherichia coli* isolates from adult and adolescent females in the United States From 2011 to 2019: Rising ESBL strains and impact on patient management. *Clinical Infectious Diseases*. 2021;73(11):1992-9.
  26. Kabugo D, Kizito, S., Dave DA, Kiwanuka AG, Nabimba R, Namunana S, K, R, Achan B, CN, F. Factors associated with community-acquired urinary tract infections among adults attending assessment centre, Mulago Hospital Uganda. *Afr Health Sci*. 2016;16:1131–42.
  27. Martin O, Adamu AA, Tibyangye J, Nyabayo J, Maniga E, Wampande C, K, D, Ezera A, B, J. Prevalence of bacterial urinary tract infections and associated factors among patients attending hospitals in Bushenyi District, Uganda. *International Journal of Microbiology*. 2019;2019.
  28. Biran A, Schmidt WP, Varadharajan KS, Rajaraman D, Kumar R, Greenland K, Gopalan B, Aunger R, Curtis VI. Effect of a behaviour-change intervention on handwashing with soap in India (SuperAmma): A cluster-randomised trial. *Lancet Glob Health*. 2014;2:45-54.
  29. Kasango SD, Lutoti S, Wewedru I, Aboce E, Angol DC. Prevalence and antimicrobial susceptibility pattern of extended spectrum beta lactamase producers in gram-negative urine isolates at MBN clinical laboratories, Kampala Uganda. *Archives of Microbiology & Immunology*. 2018;2:042-52.
  30. Kateregga JN, Kantume R, Atuhaire C, Lubowa MN, Ndukui JG. Phenotypic expression and prevalence of ESBL-producing Enterobacteriaceae in samples collected from patients in various wards of Mulago Hospital, Uganda. *BMC pharmacology and Toxicology*. 2015;16(14.).
  31. Abubakar M, Josseline N, Peruth M, Duncan O, Byaruhanga A, Ampaire L. Prevalence of extended spectrum beta-lactamase producing bacteria (ESBL) in patients presenting with urinary tract infections (UTIs) at a peri-Urban hospital, Uganda; 2020.
  32. Adler A, Katz DE, Marchaim D. The continuing plague of extended-spectrum  $\beta$ -lactamase-producing Enterobacteriaceae infections. *Infectious Disease Clinics*. 2016;30:347-375.
  33. Bitew A, Abebaw Y, Bekele D, Mihret A. Prevalence of bacterial vaginosis and associated risk factors among women complaining of genital tract infection. *International journal of microbiology*. 2017;2017(1):4919404.
  34. Anejo-Okopi JA, Okojoku OJ, Ramyil SMC, Bakwet PB, Okechalu J, Agada G, et al. Bacterial and antibiotic susceptibility pattern of urinary tract infection isolated from asymptomatic and symptomatic diabetic patients attending tertiary hospital in Jos, Nigeria. *Trends Med*. 2017; 17(1):1-5.
  35. Hemmati M, Vaziri S, Afsharian M, Mansouri F, Zamanian MH, Fereshteh S, et al. Molecular Investigation of Extended-Spectrum  $\beta$ -Lactamase and Patterns of Antibiotic Resistance in Enterobacter cloacae Isolates from Teaching Hospitals

- in Kermanshah, Iran. Journal of Clinical & Diagnostic Research. 2019;13(9).
36. Magale H, Kassim I, Odera S, Omolo M, Jaoko W, Jolly P. Antibiotic susceptibility of organisms causing urinary tract infection in patients presenting at Kenyatta national hospital, Nairobi. East African medical journal. 2015;92(7):333-7.
  37. Fatima S, Muhammad IN, Khan MN, Jamil S. Phenotypic expression and prevalence of multi drug resistant extended spectrum beta-lactamase producing Escherichia coli and Klebsiella pneumoniae in Karachi, Pakistan. Pak J Pharm Sci. 2018; 31(4):1379-84.
  38. Jena J, Sahoo RK, Debata NK, Subudhi EJB. Prevalence of TEM, SHV, and CTX-M genes of extended-spectrum  $\beta$ -lactamase-producing Escherichia coli strains isolated from urinary tract infections in adults. 3 Biotech. 2017; 7(4):244.
  39. Ogefere HO, Iriah SE, Ibadin EEJUM. Detection of SHV and TEM-type extended spectrum  $\beta$ -lactamase in bacterial isolates recovered from clinical samples of patients attending military hospitals. 2019; 38,(3).
  40. Seyedjavadi SS, Goudarzi M, Sabzehali FJJOAD. Relation between blaTEM, blaSHV and blaCTX-M genes and acute urinary tract infections. Journal of Acute Disease. 2016;5(1):71-6.
  41. Ahmed YMA, Mohammed EL, Imam M, Khalid KE, Mohamed BA, Elhag KMJBRJOM. Phenotype and genotype of extendedspectrum-beta-lactamases in Sudanese patients with UTI. 2014;3 (1-5).
  42. Hussein AA, Kassaza K, Mwesigye J, Mwamibi B, Kabanda T, Bazira J. Prevalence and genotypic characterization of extended spectrum beta-lactamase uropathogens isolated from refugees with urinary tract infections in nakivale refugee settlement camp, Southwestern Uganda. medRxiv. 2022:2022.04. 29.22274464.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/125158>