



Assessment of Maternal Mortality and Contributing Factors in Some Rural Communities of Bayelsa State, Nigeria

Onokpite Emmanuel ^a, Solomon M. Uvoh ^{b*},
Tonkiri Ayakeme ^c, Kemmer Bralate ^d
and Blessing L Dum-Awara ^e

^a Department of Anaesthesia and Intensive Care, Faculty of Clinical Medicine, College of Health Sciences, Delta State University, Abraka Delta State, Nigeria.

^b Department of Human Physiology, Faculty of Basic Medical Sciences, College of Health Sciences, University of Port Harcourt, Rivers State, Nigeria.

^c Department of Biochemistry, Faculty of Basic Medical Sciences, College of Health Sciences University of Port Harcourt River State, Nigeria.

^d Department of Community Health, Bayelsa state College of Health Sciences, Ogbia Town, Nigeria.

^e Department of Human Physiology, Faculty of Basic Medical Sciences, College of Health Sciences, Madonna University, Elele Campus, River State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126898>

Original Research Article

Received: 12/09/2024
Accepted: 15/11/2024
Published: 20/11/2024

*Corresponding author: E-mail: solomonu31@gmail.com;

Cite as: Emmanuel, Onokpite, Solomon M. Uvoh, Tonkiri Ayakeme, Kemmer Bralate, and Blessing L Dum-Awara. 2024. "Assessment of Maternal Mortality and Contributing Factors in Some Rural Communities of Bayelsa State, Nigeria". *International Journal of Research and Reports in Gynaecology* 7 (1):136-42. <https://journalijrrgy.com/index.php/IJRRGY/article/view/107>.

ABSTRACT

Aim: This study assess the implicated causes of maternal mortality in some rural communities in Bayelsa state. Communities visited having health facilities with existing records of maternal mortality from 2018-2023 include both public and private health / hospitals and TBAs centres within Ogbia LGA. Using a purposive sampling technique. Materials: The data were derived from a sproform used to record the date of maternal death during or after pregnancy within the stipulated years of study. The TBAs were asked orally the number of deaths that have occurred in their centres within the previous years.

Results: The results from this study showed a highest peak of 43.48% prevalence rate of maternal mortality after 42 days of delivery in 2019, followed by 41.43% in 2018 and 29.01% in 2023. However maternal mortality during delivery was higher in 2021 (38.78%), and 2020 (37.21%) compared with 2019 (15.22%) and 2023 of 30.69%. The main causes of maternal death observed from this study was the development of anemia during pregnancy and uncontrolled bleeding within the study years considered. Other factors contributing to maternal mortality identified were delay in seeking medical care and inadequate access to quality health care. Errors from TBAs was also a factor in 2018(23.05%), 2020(20.52%), 2022(18.32%) compared with 2023 (16.04%).

Conclusion: Findings from this study indicate a lack of awareness to the importance of engaging regularly in healthy balanced dietary foods intake during pregnancy, delay in seeking health care at the appropriate time and errors from inexperienced TBAs was the foremost cause of maternal mortality. We therefore recommend that the government should create an awareness campaign strategy to inform rural dwellers about the importance of early registration for safe delivery in health facilities located in their respective communities from the onset of pregnancy and after delivery.

Keywords: Care; health; maternal; mortality; records.

1. INTRODUCTION

“Maternal death refer to a woman’s death during pregnancy or within 42 days after abortion or delivery, irrespective of the duration and place of the pregnancy, from any cause related to or aggravated by the pregnancy or its management but not from accidental or unintentional causes” (Ujah et al., 2005; Khama et al., 2006; Owolobi et al., 2019; Nivin, 2019; Oghenetega et al., 2020). “The growing concern on improving reproductive health at the global level has created a demand for research especially in the area of maternal health. Maternal health which is the physical wellbeing of a woman during pregnancy, childbirth and postpartum period. However reports from WHO in 2020 showed an estimated death of 1047 in every 100,000 in every livebirth” (Alkema et al., 2015; Fadeji, 2017; Solomon et al., 2021; WHO, 2021; Hamilton et al., 2023)

“It is within this conceptual framework that the millennium development goal target 5A, calls for a reduction in maternal mortality ratio by three-quarters by 2015. At its present rate, the world will fall short of the target for maternal mortality reduction because the data collated suggest that to reach the target, the global maternal mortality rate would have had to be reduced by an average of 5.5. % between 1990 and 2015”

(Khama, 2006; Ogunjuyigbe and Liasu, 2007; NDHS).

Nigeria constitute only two percent of the world’s population but account for over 10% of the world maternal deaths, and ranks second globally only to (Okonofua, 2007). The status of maternal health is poor in Nigeria, defined by maternal mortality of 59,000 per annum due to pregnancy related causes. This has identified as the leading cause or determinant of death among women of reproductive age in Nigeria. Although opinion differ on the determinants of maternal mortality. Herfon (2006), noted that the causes of maternal mortality is an outcome of nexus interaction of a variety of factors namely; the distant factors (socio-economic, income level and illiteracy act through the proximate or intermediate factors (health and reproductive behavior, access to health services) and in turn influence outcome of pregnancy complication mortality. further identified other factors responsible for maternal mortality as socio-cultural factors which includes; traditional practices, norms, beliefs, education and religion. Several attempts have been made in the past aimed at reducing maternal mortality in Nigeria, such attempts, especially by the federal and state governments; have generally not proved very successful in achieving the desired results. Some promising results however

have recently begun to be recorded through some policy initiatives by a few state governments (Ogunjuyigbe and Liasu, 2013; Fabio et al., 2015, 2016).

The introduction of the safe motherhood program in 1995, midwife service scheme in 2011 and subsidy reinvestment program (SURE-P) in 2012 introduced a range of intervention which included antenatal, postnatal care, family planning prevention and management of unsafe abortions, and health education but still maternal mortality rate has not been encouraging over the years and improvement are so low. Maternal Mortality remains a key issue affecting women of reproductive age across the African Region. Despite the global decline in the maternal mortality ratio to 34.2% between 2000 and 2020, MMR continue to be a disaster in the Africa region. With more than two-thirds (69%) of maternal deaths occurring in the African Region.

From recent estimates, the number of deaths each year from maternal causes worldwide decreased from 536,000 in 2005 to an estimated 358,000 in 2008 and 273,500 in 2011. "For every woman that dies approximately 20 more suffers injuries and infections associated with disabilities in pregnancy or childbirth" (Hamilton, 1949; Jochen et al., 2011; Grindheim, 2015) Solomon et al., 2022). The situation is even more alarming in Nigeria for example, in the year 2000, the maternal mortality ratio per 100,000 live births was 800 compared to 540 for Ghana and 240 for South Africa.

Consequently, the chances of a Nigerian woman dying from Reproductive health disorders and complications was put at 1 in 10 in 2002 Population Reference Bureau, 1 in 5 in 2005, and 1 in 23 in 2008 placing the Nigerian women at far greater risk than her counterpart in developed world where the risk is estimated to be 1 in 17,800 and 1 in 10,000 in countries such as the republic of Ireland and Singapore respectively (Charles et al.,1990,1998; World Bank, 2011;Eric et al.,2013; Ovuakporaye et al.,2016;Blessing et al., 2023). Some of the implications of these estimates are the depletion of the country's workforce and the overall stifling of rapid development.

There are no studies to inform the populace in the current study area, hence, this study focuses on maternal mortality and causes in Ogbia L.G.A in Bayelsa State.

2. MATERIALS AND METHODS

2.1 Study Design

Retrospective research design was employed in this study

2.2 Area of the Study

The study area lies within the saltwater and freshwater swamp geographic units of the Niger Delta.It includes Otuabagi, Oloibiri, Otuakeme, kolo, Otuasega, Elebele, otueke, Emeyal , Opume, and TBAs centres. The inhabitants of Ogbia land are mainly fishermen and farmers.

2.3 Study Population

The population of this study consist of all the existing records of maternal mortality cases from the health facilities (both private and public) including TBAs centers within the study area.

2.4 Sample Size

The sample size include all the existing records of maternal mortality in the various health facilities from 2018 to 2023.

2.5 Sampling Technique

A purposive sampling technique was used to obtain samples from 2018 - 2023.

2.6 Nature/Source of Data

The study derive its data from secondary source through the use of a preform used to record all date retrieved from the records

2.7 Instrument for Data Collection

The instrument for data collection was sproform used to record all data retrieved from the records and direct oral findings from TBAs.

2.7.1 Data collection

The researchers recruited some assistants during the collection process to carefully check files and record of all maternal mortality cases and causes from the stipulated years

2.7.2 Data analysis

Data analysis was done using SPSS software Version 24.0 with results presented in tables, charts expressed as frequencies and percentages.

3. RESULTS

Table 1 shows the prevalence of maternal mortality; in 2018 with 41.43% of maternal mortality within 42 days of delivery, in 2019, 43.48% of maternal mortality within 42 days of delivery, in 2020, 37.21% of maternal mortality during childbirth, in 2021, 38.78% of maternal mortality during childbirth, in 2022 39.02% of maternal mortality while pregnant, in 2023 40.32% of maternal mortality while pregnant.

Table 2 showing causes of maternal mortality; in 2018, 37.69% agreed on anemia during pregnancy, in 2019, 37.14% agreed on anemia during pregnancy, in 2020, 43.97% also agreed on anemia during pregnancy, in 2021, 36.59% agreed on anemia during pregnancy, in 2022, 42.78% agreed on anemia during pregnancy, in 2023, 42.71% also consented to anemia during pregnancy being the cause of maternal mortality.

Table 3 showing the contributory factors to maternal mortality; in 2018, 26.29% consented to delay in seeking medical care, in 2019, 22.01% believed in delay to seek care, in 2020, 20.53% agreed on errors from traditional birth attendants, in 2021, 16.37% believed on errors from

traditional birth attendants, in 2022, 19.37% agreed on delay in seeking care, in 2023, 16.04% agreed that errors from traditional birth attendants was a contributory factor to maternal mortality.

4. DISCUSSION

Concerning the causes of Maternal mortality, the study showed that bleeding after delivery (23.81%) was the leading cause of mortality in 2018, pre-eclampsia or eclampsia during pregnancy (13.49%) was the highest percentage in 2019, bleeding after delivery (19.27%) in 2020, bleeding after delivery (21.95%) in 2021 while anemia during pregnancy result in 42.78% as the major causes of mortality in 2022. However anemia account for mortality peak rate of (42.71%) in 2023.

Concerning contributing factors to maternal mortality, this study observed that delay in seeking care was the most causes of maternal mortality with 26.29%, while results from 2019 shows that delay in seeking care was at 22.01% maternal mortality rate, and errors from traditional birth attendant (20.53%) was the highest causes of maternal mortality in 2020.

Table 1. Prevalence of maternal mortality in 2018-2023

Mortality	2018	2019	2020	2021	2022	2023
	(n=70)	(n=46)	(n=43)	(n=49)	(n=41)	n =62
	F (%)	F (%)	F (%)	F (%)	F (%)	F (%)
During pregnancy	25(35.71)	19(41.30)	12(27.91)	17(34.69)	16(39.02)	25(40.32)
During child birth	16(22.86)	7(15.22)	16(37.21)	19(38.78%)	10(24.39)	19(30.65)
Within 42 days of delivery	29(41.43)	20(43.48)	15((34.88)	13(26.53)	15(36.59)	18(29.03)
otal (%)	100	100	100	100	100	100

Table 2. Causes of maternal mortality in the study area

Variables	2018	2019	2020	2021	2022	2023
	(n=252)	(n=175)	(n=166)	(n=164)	(n=173)	(n=192)
Bleeding after delivery	60(23.81)	42(24)	32(19.28)	36(21.95)	23(13.29)	41(21.35)
Pre-eclampsia/ Eclampsia during pregnancy	34(13.49)	27(15.43)	11(6.63)	17(10.37)	25(14.45)	21(10.94)
Anemia during pregnancy	95(37.69)	65(37.14)	73(43.97)	60(36.59)	74(42.78)	82(42.71)
Poor Maternal health care	43(17.07)	26(14.86)	28(16.87)	26(15.85)	29(16.76)	23(11.98)
Educational status	20(7.94)	15(8.57)	22(13.25)	25(15.24)	22(12.72)	25(13.02)

Source: Field survey, 2024

Table 3. Factors contributing to maternal mortality in the study area

Variable	2018 n= 308		2019 n= 209		2020 n=190		2021 n= 171		2022 n=191		2023 n=212	
	F	%	F	%	F	%	F	%	F	%	F	%
Inadequate human resource for health	20	6.49	23	11.01	16	8.42	13	7.60	23	12.04	19	8.96
Delay in seeking care	81	26.29	46	22.01	36	18.95	21	12.28	37	19.37	28	13.21
Lack of access to quality care	29	9.42	17	8.13	20	10.53	18	10.53	19	9.95	27	12.74
Inadequate equipment	20	6.49	17	8.13	18	9.47	19	11.11	14	7.33	23	10.85
Lack of ambulance transportation	17	5.52	8	3.83	12	6.32	16	9.36	13	6.81	16	7.54
Delay in referral services	37	12.02	21	10.05	14	7.37	26	15.21	21	10.9	22	10.38
Errors from traditional birth attendants	71	23.05	40	19.14	39	20.53	28	16.37	35	18.32	34	16.04
Errors from prayer or mission houses	23	7.47	21	10.05	18	9.47	15	8.77	14	7.33	22	10.38
Diarrheal related death e.g. poor access to water, sanitation and hygiene	10	3.25	16	7.66	17	8.95	15	8.77	15	7.85	21	9.91

Source: Field survey, 2024

However 16.37% mortality was due to errors from traditional birth attendant in 2021 and 19.37% as the leading cause of maternal mortality in 2022 while 16.04% was observed from errors from traditional birth attendants in 2023.

5. CONCLUSION

Maternal mortality remains a challenge in global health, with well-known disparities across countries. However this study showed errors from prayers without doing the needful medically at the appropriate time, lack of ambulance to transport pregnant women in critical condition, and poor management of pregnancy induced anemia are some of the causes of maternal mortality identify among pregnant women in Ogbia local government area of Bayelsa state.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Institutional ethical approval was obtained from the Research Ethics Committee and permission from the management of the health facilities visited for easy access to the medical records of maternal morbidity was also sorted.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Black RE, Laxminarayan R, Temmerman M, (2016). Reproductive, Maternal, Newborn, and Child Health: Disease Control Priorities, *Publication of international bank for reconstruction and development / the World Bank*; 20(16) 5-20. doi: 10.1596/978-1-4648-0348-2_
- Charles RB, Frank WL, Williams NPH, Douglas WL, Rogers PS, Barbara MB (1998,1990). *Obstetrics and gynecology* (3rd edition) Williams and Wilkins ISBN 0-683-30391-0.
- Eric DL, Joshua R, Chair Whitney AS, Roger AB, Catherine Proeza (2013). Depressed pacemaker activities of sino-arterial node

myocytes contribute to age dependent decline in maximum heart rate. *Proceedings of the National Academy of Sciences of the United State of America (PNAS)*.

- Fabio Ageli, Enrica Angeli, and Paolo Verdecchia (2015). *Int. J. Mol. Sci.* 16, 18458-18473 Review Novel electrocardiographic patterns for the prediction of hypertensive disorders in pregnancy – from pathophysiology to practical implications.
- Fabio PA Lourenco AS, Hello BS, Marcos VT, Nilo B (2016). Bioaccumulation of mercury, cadmium, Zinc, Chromium, and lead in muscle, liver and spleen tissues of a large commercially valuable catfish species in Brazil. *Anas of the Brazilian Academy Sci* 88(1): 137-147.
- FilippiV, and ChouD, (2016). Levels and Causes of Maternal Mortality and Morbidity. *Journal of library books*. https://doi.org/10.1596/978-1-4648-0348-2_ch3
- Grindheim E, Toska K, Estensen MF, Rosseland LA. (2011,2015). Changes in pulmonary Function during pregnancy: a longitudinal Cohort Study *BJOG* 119:94-101.
- Hamilton HFH (1949). Cardiac output in normal pregnancy as determined by Cournand right heart catheterization techniques *Journal of obstetrics and Gynecology Br. Emp* 56,54.
- Hamilton dos PT, Job MT, Suelman BM, Prata T (2023). Maternal mortality: A matter of public Health Policies. DOI:10.4236/0j0g.2023.136088
- Herfon (2006). Health Reform Foundation of Nigeria. *Nigeria Health Review* Kembim press Ltd Ibadan pp 10-12,130,191.
- Jochen Steppan, Viachaslau M Barodka, Dan E Berkwitz, Daniel Nirhan (2011). Vascular stiffness and increased pulse pressure in the ageing cardiovascular system. *Journal of cardiology Research and practice* (2):263585.
- Khama OR, John O, Philip M (2006). Disease and mortality in sub-Sahara Africa 2nd ed Washington DC; The International Bank for Reconstruction and Development/ the World Bank Chapter 16.
- Musarandega R, Nyakura M, Machezano R, Pattinson R, and Munjanja SP, (2021). Causes of maternal mortality in Sub-Saharan Africa: A systematic review of studies published from 2015 to 2020. *Journal of global health.* 12(4).12-23. DOI: 10.7189/jogh.11.04048

- Nigeria Demographic and Health Survey (NDHS). Latest edition. Nivin Todd MD (2019). Potential complication: Gestational hypertension-Web MD.
- Oghenetega Onome, Ana G, Okunlola M, Ogengbede O(2020). Oil Spills, Gas Flaring and adverse pregnancy outcomes. A systematic review. Journal of obstetrics and gynecology, 10,187-199.
- Ogunjuyigbe and Liasu (2013). Impact of child health on economic growth: New evidence based on granger on-casualty test.
- Okonofua E (2007). New Research Findings on Adolescent Reproductive Health in Nigeria. African Journal of Reproductive Health: 17-19.
- Oluwatoyin OA, Akpoghomeh OS, (2019). Analysis of the Temporal and Seasonal Patterns of Maternal Mortality Ratio in yenagoa, Bayelsa state of Nigeria. *International journal of research in geography (IJRG)* 5(2). 13-18. ISSN 2454-8685 <http://dx.doi.org/10.20431/2454-8685.0502002> www.arcjournals.org
- Ovuakporaye Sol, Igweh CJ, Aloamaka CP (2016). Imp-act of gas flaring on cardiopulmonary parameters of residents in gas flowing communities in Niger Delta Nigeria. *British Journal of Medicine and Medical research* 15 (6):1-13, (BJMMR).
- Blessing LD, Solomon MU, Emily Kiridi GE, Onokpite E (2023). Oxidative stress markers and lipid profile among female residents in oil impacted communities of River State, Nigeria. *International Journal of Clinical Studies & Medical Case Reports*.Vol.28 (1):1-7
- Sageer R, Kongnyuy E, AdebimpeWO, Omosehina O, Ogunsola EA,&Sanni B(2019). Causes and contributory factors of maternal mortality: evidence from maternal and perinatal death surveillance and response in Ogun state, Southwest Nigeria. *Biomedical journal of pregnancy and childbirth* 19(63): 892-974.
- Solomon MU, Tonkiri A, Onokpite E, Blessing DL (2022). Nephrotoxicity among adult male residents exposed to gas flares in Bayelsa state Nigeria. *Quest Journals Journal of Medical and Dental Science Research Volume 9~ Issue 4 pp.: 59-63* ISSN (Online) : 2394-076X ISSN (Print):2394-0751 www.questjournals.org
- Solomon MU, Charles NN, Emily KGE (2021).Blood serum lead and cadmium level among pregnant women in gas flaring communities in Bayelsa State, Nigeria.
- Solomon MU, Arthur NC, Obia B (2021).Impact of gas flares on anthropometric indices of pregnant and non-pregnant women in selected gas flaring communities in Bayelsa state, Nigeria. *GPH International Journal of Research in Biological and Medicine Science* 4(9): 15-21.
- Solomon MU, Arthur NC, Kiridi Emily GE, Charles NN (2021).Toxic air and respiratory indices among pregnant women in Bayelsa state, Nigeria. *Spectacular Journal of Research in Medicine and Medical Sciences* 2(5):099-103.
- Solomon MU, Azibalua AA, Kiridi Emily GE, Alagoa Bibi-welson E, Blessing DL(2021).Total Serum Cholesterol and Selected electrolytes during pregnancy among women residents in yenagoa metropolis in Bayelsa State, Nigeria.
- Ujah AO, Aisien OA, Mutahir JT, Vanderagt DJ,Glew RH et al (2005).Socio- Cultural Factors of Gender Roles in Women Health Care Utilization in Southwest Nigeria. *Journal of Obstetrics and Gynecology* 13(6).
- World Health Organization. (2021). Maternal mortality: The agency of a systemic and multi sectoral approach in mitigating maternal deaths in Africa.Available at: (IAHO) aho.afro.who.int
- United Nations Population Fund (UNFPA) (2023). Report on Maternal Health in Nigeria:Ending preventable maternal deaths By 2025, the reduction of preventable maternal deaths has accelerated.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/126898>