



Ulcerative Colitis in Children at Campus Teaching Hospital of Lome-Togo: A Report of Two Cases

Laté Mawuli Lawson-Ananissoh ^{a*}, Rafiou El-Hadji Yakoubou ^b,
Koffi Mawuse Guédénon ^c, Debehoma Venceslas Redah ^a,
Lidawu Roland-Moise Kogoe ^a, Oumboma Bouglouga ^b and Aklesso Bagny ^a

^a Service of Hepatology and Gastroenterology, Campus Teaching Hospital, Lome, Togo.

^b Service of Hepatology and Gastroenterology, Kara Teaching Hospital, Kara, Togo.

^c Pediatric Service, Sylvanus Olympio Teaching Hospital, Lome, Togo.

Authors' contributions

. This work was carried out in collaboration among all authors. Author LMLA conceptualized the study, performed literature review and wrote the initial draft. Author REHY managed the patient, assisted with the initial draft and performed literature search. Author KMG managed the analyses of the study. All authors read and approved the final manuscript

Article Information

Editor(s):

(1) Dr. P. Veera Muthumari, V.V.Vanniaperumal College for Women, India.

Reviewers:

(1) Ulrica Lovén Wickman, Linnaeus University, Sweden.

(2) Fatemeh Ghani Dehkordi, Shiraz University of Medical Sciences, Iran.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/81958>

Original Research Article

Received 20 October 2021

Accepted 27 December 2021

Published 29 December 2021

ABSTRACT

Introduction: Ulcerative colitis is one of the chronic inflammatory bowel diseases which, together with Crohn's disease. We aimed to report the first cases of ulcerative colitis in two children in Togo and describe their diagnostic and therapeutic course.

Result: The first case concerned a 30-month-old infant brought in by his parents was admitted for glairo-sanguinous diarrhea that had been evolving for 13 months before the consultation. The parents consulted general practitioners and pediatricians on several occasions without success. Total colonoscopy with biopsies and their anatomopathological examination were in favor of ulcerative colitis. The blood test for ASCA was negative and for pANCA positive. The second case concerned a 12-year-old teenager accompanied by his mother and was seen in consultation for daily rectal bleeding evolving since the age of about 7 years, which sprays the toilet bowl regardless of the consistency of the stools. Total colonoscopy with biopsies and their anatomopathological examination noted a rectal mucosa with slight chronic inflammatory changes

*Corresponding author: Email: lawsonprosper@yahoo.fr;

without specificity. The blood test for ASCA was negative and that of pANCA positive. The two patients were put on mesalamine with a good evolution under treatment.

Conclusion: UC should be rapidly evoked in children and infants presenting symptoms such as rectal bleeding and chronic diarrhea in order to ensure early diagnosis and care.

Keywords: Children; inflammatory bowel disease; Togo; ulcerative colitis.

1. INTRODUCTION

Ulcerative colitis (UC) is a chronic inflammation of the digestive tract exclusively affecting the rectum and colon [1]. It is one of the chronic inflammatory bowel diseases (IBD) which, together with Crohn's disease, constitute a public health problem in industrialized countries, where they mainly affect young adults [2]. However, these diseases are clearly on the rise in developing countries and are increasingly diagnosed in young subjects and children from the age of 15 [3]. UC is described as rare in Africa compared to the West [4]. In Togo, Bagny et al reported a case of IBD in a 14-year-old teenager but it concerned Crohn's disease [5]. The purposes of this work were to report the first cases of UC in two children in Togo and to describe their diagnostic and therapeutic course.

2. RESULTS

2.1 Observation 1

A 30-month-old infant brought in by his parents was admitted for glairo-sanguinous diarrhea that had been evolving for 13 months before the consultation at a rate of 6 to 8 stools per day. He is the second child of the family whose eldest is 5 years old alive and well. He had no fever or vomiting. There was a progressive deterioration of the general condition with a marked anemia (hemoglobin level at 6.8 g/dl) that required a blood transfusion in a health facility. The parents consulted general practitioners and pediatricians on several occasions. He was regularly put on fluconazole and metronidazole without success. On examination, the child was very irritable and weighed 9.2 kg, not febrile and in poor general condition. Total colonoscopy performed under sedation down to the caecal floor had noted a very erythematous recto-colic mucosa as a whole, fragile and bleeding easily on contact; with disseminated ulcerations; multiple staged biopsies were performed. The anatomopathological examination of the biopsies noted: the colonic mucosa shows a weakened and sometimes detached lining. The chorion is thickened by a polymorphic inflammatory

infiltrate of marked density; the glands are sometimes elongated and bordered with a partially depleted mucosal lining; without dysplasia; cryptic abscesses are visible; congestion and edema are moderate without specific lesions; this was in favor of ulcerative colitis. The blood test for ASCA was negative and for pANCA positive. The blood count showed a hemoglobin level of 9.2 g / dl, 10,000 leukocytes with 6,200 polymorphonuclear neutrophils; the C reactive protein was 17 mg / l; the sedimentation rate was 77 mm in the first minute; serum creatinine at 2 mg / l; the liver function test and blood ionogram were normal. The patient was put on mesalamine (granules) 500 mg daily. One month after the start of treatment, there was a persistence of diarrhoea at the rate of 3 to 5 stools per day always glairo-sanguinous; it was thus associated with corticosteroid therapy containing prednisone 10 mg per day for 14 days. At the end of this association with corticosteroid therapy, there was a reduction in the number of stools to 2 per day, not glairo-sanguinous, with a normal appearance; a better clinical condition with a weight of 10.5 kg; a non-irritable child; mesalamine was continued after stopping corticosteroid therapy. Six months after the start of treatment, 2 stools per day with normal appearance were noted.

2.2 Observation 2

A 12-year-old teenager accompanied by his mother was seen in consultation for daily rectal bleeding evolving since the age of about 7 years, which sprays the toilet bowl regardless of the consistency of the stools without any notion of mucus or deterioration of the general condition; neither abdominal pain nor proctalgia nor diarrhea nor constipation. He is the 5th in a family of 5 children with living healthy siblings; no particular history in the parents. The patient has received several modern (metronidazole, albendazole, ciprofloxacin) and traditional treatments without success. On examination of this teenager in good general condition and weighing 42 kg, a healthy anal margin without fissure was noted; total colonoscopy performed

up to the caecal floor noted an inflammatory zone 5 to 6 cm from the anal margin; circumferential erythematous of about 2 cm in height, bleeding easily on contact and biopsied; the rest of the colonic mucosa was of normal appearance. The anatomopathological examination noted a rectal mucosa with slight chronic inflammatory changes without specificity. The blood test for ASCA was negative and that of pANCA positive. The biological assessment noted a creatinine level of 5 mg/l; the hemoglobin level was 11g/dl microcytic hypochromic; the C-reactive protein was 5 mg/l. A mesalamine-based treatment (suppository) was initiated at a rate of 1g x 2 per day. An improvement in rectal bleeding was noted after 1 month of treatment. At 6 months after the start of mesalamine treatment, no more rectal bleeding was noted.

3. DISCUSSION

These two clinical cases studied in our work are the first cases reported in Togo in children. Bagny et al had described a case of Crohn's disease in an adolescent girl. in sub-Saharan Africa, we have not found cases of ulcerative colitis described in children to our knowledge. IBD and in particular UC are very rare conditions in children. They are often increasingly diagnosed in young subjects and teenagers [6]. Infants and toddlers most often present with several digestive signs related to other causes, particularly infectious ones. It is the persistence of symptoms after adolescence that draws the attention of practitioners to IBD. Our two patients are male. A male predominance of UC is noted in the literature but the opposite has been observed in children and remains unexplained [7]. The pathophysiology of UC or IBD in general involves many environmental and genetic risk factors. Environmental factors are not well known but seem to be associated with the modern western lifestyle [8]. There is a familial predisposition for UC although much lower than for Crohn's disease. It is a disease that can affect several members of the same family with so-called "susceptibility" genes. These genes involved in the genetic predisposition to IBD are very numerous (greater than 160) [9]. The percentage of familial forms of IBD varies between 5 and 20% in the literature [10]. The diagnosis of ulcerative colitis is often delayed and difficult and is based on a number of clinical and histological arguments. The delay between the onset of symptoms and the diagnosis is more than one year in our patients. This delay is significantly longer than that observed in France for the

diagnosis of IBD, whose median is about 3 months [11]. This long delay in diagnostic in our context noted by other authors [5,12] could be explained by the scarcity of gastroenterologists (8 for the whole country) and gastro-pediatricians (none in the country). Patients repeatedly consult general practitioners who do not perform endoscopic explorations and often dismiss the symptoms as simple enteritis. Total colonoscopy with biopsies for histological examination is essential for diagnosis. The presence of a biological inflammatory syndrome and perinuclear anti-neutrophil antibodies (pANCA) are also important elements that help make the diagnosis of UC. Therapeutically, the use of traditional drugs and antibiotics by our patients prior to diagnosis is a factor that could worsen the disease with a disorganization of the intestinal microbiota. Treatment of UC involves medications that regulate the functioning of the patient's immune system, anti-inflammatory drugs and immunosuppressants. Anti-TNF drugs are also used in children with indications and dosages comparable to those of adults [13]. Another pediatric specificity is the use of nutritional assistance, mainly in the form of cyclic enteral nutrition, which is as effective as corticosteroids [14] but anti-TNFs and nutritional assistance are not available locally. Our two patients benefited from mesalamine with a good clinical evolution and amendment of symptoms. This treatment must be maintained over the long term, hence the need for therapeutic education involving the parents.

4. CONCLUSION

A delay in diagnosis of ulcerative colitis in our context was noted. It may be due to the delay in carrying out the total colonoscopy with biopsies and their anatomopathological examination. UC should be rapidly evoked in children and infants presenting symptoms such as rectal bleeding and chronic diarrhea in order to ensure early diagnosis and care.

CONSENT

Written informed consent was obtained from the parents of the children included in the study.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENTS

We appreciate the cooperation of the parents of the children who agreed to participate in the study. Our thanks go to the management of the Campus Teaching Hospital for the support with ideal environment to perform the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Torres J, Billioud V, Sachar DB, Peyrin-Biroulet L, Colombel JF. Ulcerative colitis as a progressive disease: the forgotten evidence. *Inflamm Bowel Dis.* 2012; 18:1356-63.
2. Loftus E.V. Clinical epidemiology of inflammatory bowel disease: incidence, prevalence and environmental influences. *Gastroenterology.* 2004;126:1504-17.
3. Moline F., Gower-Rousseau C., Yzet T. et al. Opposite evolution in incidence of Crohn's disease and ulcerative colitis in Northern France (1988-1999). *Gut.* 2004; 53:843-8.
4. Alese OB, Irabor DO. Pyoderma gangrenosum and ulcerative colitis in the tropics. *Rev Soc. Bras. Med. Trop.* 2008; 41:6647.
5. 5-Bagny A, Akolly DAE, Kogoe LRM, Djadou KE. A girl with Crohn's disease in a tertiary care hospital of Togo: A Case Report. *Asian Journal of Research and Reports in Gastroenterology.* 2021;5(1): 28-31
6. Jean-Frédéric C, Gwenola VM, Antoine C, Gower-Rousseau C, et al. Jean-louis S. Épidémiologie et facteurs de risque des maladies inflammatoires chroniques de l'intestin. *Bull. Acad. Natle Med.* 2007; 191:1105-1123.
7. Sach SC, Khalili H, Gower-Rousseau C, Olen O, Benchimol EI, Lynge E, et al. Sex-based differences in incidence of inflammatory bowel diseases-pooled analysis of population-based studies from western countries. *Gastroenterology.* 2018; 155(4):1079-89.
8. POST U. Hugot JP, Viala J. Physiopathologie de la maladie de Crohn: l'indispensable pour le clinicien. 2015;181-8.
9. Jostins L, Ripke S, Weersma RK, et al. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature.* 2012;491(7422):119-24.
10. Ahmad T, Tamboli CP, Jewell D, Colombel JF. Clinical relevance of advances in genetics and pharmacogenetics of IBD. *Gastroenterology.* 2004;126:1533-49.
11. Gower-Rousseau C, Leryer A, Génin M, et al. Epidémiologie descriptive et évolution dans le temps et l'espace de l'incidence des maladies inflammatoires chroniques intestinales dans le nord-ouest de la France (1988-2014). *Bull Epidémiol Hebd.* 2019;(13):228-36.
12. Bougouma A, Sombié R, Goumbri-Lompo O, Ido-Da TRC, Zoure N, Darankoum D et al. La rectocolite hémorragique chez le noir africain. Analyse de 20 cas au Burkina Faso. *J. Afr. Hépatol. Gastroentérol.* 2010; 4:147-151.
13. Hyams J, Crandall W, Kugathasan S, Griffiths A, Olson A, Johans J, et al. Reach Study Group. Induction and maintenance infliximab therapy for the treatment of moderate-to-severe Crohn's disease in children. *Gastroenterology.* 2007;132(3): 863-73
14. Day AS, Whitten KE, Sidler M, Lemberg DA. Systematic review: nutritional therapy in paediatric Crohn's disease. *Aliment Pharmacol Ther.* 2008;27(4):293-307.

© 2021 Lawson-Ananissoh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/81958>