



## **Detection of Dichlorvos Residue in Cowpea Grains, Six Months after Application Using High Performance Liquid Chromatography**

**S. R. Yusuf<sup>1\*</sup>, S. H. Lawan<sup>1</sup>, B. S. Wudil<sup>1</sup> and H. Sule<sup>1</sup>**

<sup>1</sup>*Department Crop Protection, Bayero University, P.M.B. 3011, Kano, Nigeria.*

### **Authors' contributions**

*This work was carried out in collaboration between all authors. Author SHL designed the study and conducted the research work. Author SRY, wrote the first draft of the manuscript and made conceptual contributions, corrections and objective criticisms, he was assisted by authors BSW and HS. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/ARJA/2017/37992

#### Editor(s):

(1) M. Yuvaraj, Assistant Professor, Department of Soil Science and Agricultural Chemistry, Adhiparasakthi Agricultural College, Tamil Nadu, India.

#### Reviewers:

(1) Cláudia Helena Pastor Piscato, Brazil.

(2) Aba-Toumnou Lucie, University of Bangui, Central African Republic.

Complete Peer review History: <http://www.sciencedomain.org/review-history/22570>

**Short Communication**

**Received 6<sup>th</sup> November 2017**  
**Accepted 27<sup>th</sup> November 2017**  
**Published 2<sup>nd</sup> January 2018**

### **ABSTRACT**

In Kano State Nigeria, especially at the famous Dawanau grains market, dichlorvos (DDVP: 2, 2-dichlorovinyl dimethyl phosphate), which is a synthetic organophosphate insecticide is being widely used against stored insect pests of cowpea grains by traders and merchants alike. High Performance Liquid Chromatography (HPLC) screening using commercially available brand of such an organophosphate insecticide was carried out in a Chemistry Laboratory at Bayero University Kano in 2014. This was aimed at determining the presence or absence of the insecticide in the treated and untreated grains of cowpea before, day one and six months after application. The analysis was carried out using an Agilent 1260 Quadpump HPLC, equipped with diode array detector, autosampler and thermostatted column compartment. The result was processed using Agilent Chemstation software. Dichlorvos detection was achieved at 214 nm, while the overlaid chromatograms suggested the presence of DDVP: 2, 2-dichlorovinyl dimethyl phosphate in the test sample relative to the chromatogram of the standard commercial sample at a retention time of 2.163 and 2.283 minutes, respectively. The peak of the target compound was visibly not present in the control chromatogram at the stated retention time. In conclusion, detection of dichlorvos in grains six

\*Corresponding author: E-mail: [saniryusuf@gmail.com](mailto:saniryusuf@gmail.com);  
E-mail: [babscom68@gmail.com](mailto:babscom68@gmail.com);

months after application highlights among others, the dangers and/or disadvantages associated with the use of the chemicals for the protection of stored cowpea grains against insect pests especially by the personnel lacking requisite knowledge on the appropriate ways of handling/application of such synthetic products for both human and environmental safety.

*Keywords: Dichlorvos; organophosphate; agilent 1260 Quadpump HPLC; chromatogram; autosampler; DDVP: 2; 2-dichlorovinyl dimethyl phosphate and diode array detector.*

## 1. INTRODUCTION

Food security in sub-Saharan Africa largely depends upon improved food productivity through the use of sustainable agricultural practices and the reduction of post-harvest losses caused by pests and diseases [1, 2]. According to [3] pests and diseases pose the greatest threat to increased food production during storage and handling. Before harvest, insects cause about 15 – 100% crop damage, while storage losses in food grain range between 10 – 60%. Storage insect pests cause substantial damage to the stored grains [4]. However, [5] pointed out that to ensure high food quality and standards acceptable to the consumer, quality control, including good storage and handling practices must be observed at all stages. Managing stored grains requires the use of various techniques to ensure that the quality of the stored grains does not deteriorate over time. These measures include: the use of sanitation; storing sound, whole and dry grain; managing temperature and aeration; including the use of chemical protectants, regular sampling, and fumigation [6].

Pesticide usage in developing nations like Africa has increased dramatically in recent times with adverse effects on humans and other non-target organisms [7]. Misuse of highly toxic pesticides, coupled with a weak or a totally absent legislative framework in the use of pesticides are reasons for the high incidence of pesticide poisoning in developing countries [8]. Unintended exposure to pesticides can occur during their manufacturing, formulation and application or from environmental residues after application [9]. The use of pesticides in grains storage becomes more intensive in the Northern part of Nigeria especially at Dawanau grains market in Kano State, which is viewed as the largest grain market in West Africa. The cowpea grain farmers and traders' ignorance about pesticide toxicity led to its misuse: abusive application, inadequate usage, etc. [10].

Dichlorvos (2, 3-dichlorovinyl dimethyl phosphate) is one of the classes of insecticides

referred to as organophosphates used to control households and stored products insects. It is effective against mushroom flies, aphids, spider mites, caterpillars, thrips, and white flies in greenhouse, outdoor fruits, and vegetable crops [11,12]. Therapeutically, dichlorvos is used as a fumigant or to treat a variety of parasitic worm infections in dogs, livestock and humans. It acts against insects as a contact and stomach poison [11]. Dichlorvos pesticide self-poisoning is an important clinical problem in the developing world, and kills an estimated 200,000 people every year [13]. Concentrates of dichlorvos is mildly irritating to skin and may cause localized sweating, involuntary muscle contractions and burning sensations or actual burns [14]. When inhaled, the first effects are usually respiratory and may include bloody or runny nose, coughing, chest discomfort, difficult or short breath, and wheezing due to constriction or excess fluid in the bronchial tubes. When in contact with eyes will cause pain, bleeding, tears, pupil constriction, and blurred vision [12]. The objective of this study is to determine the presence or absence of dichlorvos in treated cowpea grains six months after storage.

## 2. MATERIALS AND METHODS

The study was carried out at the Laboratory of the Department of Pure and Industrial Chemistry, Faculty of Science, Bayero University Kano (11° 58' N, 8° 25' E) in 2014. Five kg of untreated cowpea grain (Dan Misra) was obtained from the Kano Agricultural and Rural Development Authority (KNARDA) this was divided into two i.e 4 kg which was treated with dichlorvos (4 ml/4 kg) and 1 kg (untreated control), simulating the dose of dichlorvos used by cowpea merchants at Dawanau market for the protection of cowpea grains while in storage. According to the Codex Alimentarius of the Food and Agricultural Organisations and the World Health Organisation that the mean dichlorvos concentration in cowpea ranged from 0.001 to 0.108 mg/kg<sup>-1</sup> for organochlorine pesticides, 0.002 to 0.015 mg/kg<sup>-1</sup> for organophosphorus pesticides and 0.001–0.039 mg/kg<sup>-1</sup> for pyrethroids pesticides [13].

Six months after, 50 g of the dichlorvos treated and untreated cowpea seeds were measured and each put in a 250 ml conical flask after which 100 ml of methanol was added in each flask, the treated cowpea was washed using the methanol to extract the expected dichlorvos residues from the cowpea. The mixture was then used for the screening exercise. Qualitative screening of treated and untreated cowpea samples for the detection of presence or absence of Dichlorvos insecticide was made using the high performance liquid chromatography and commercially available brand of dichlorvos was used as a guide. A 20 minutes program was carried out using water/acetonitrile gradient on Zorbax high definition 3.5  $\mu$ -C18-column (RP), at 20°C, 20  $\mu$ L was injected from a solution of 10 mg/ml in HPLC grade methanol. Analysis was run at 1 ml per minute and detection was achieved at 214 nm. The analysis was conducted with Agilent 1260 Quadpump HPLC, with diode array detector, autosampler and thermostatted column compartment. The result was processed using Agilent Chemstation software.

### 3. RESULTS

Fig. 1 reveals the presence of the target insecticide (dichlorvos) in the protected cowpea stored for the duration of six months in the store and the detection was made at 214 nm. The overlaid chromatograms suggested the presence of DDVP: 2, 2-dichlorovinyl dimethyl phosphate

in the test sample relative to the chromatogram of the standard commercial sample (Fig. 2) at a retention time of 2.163 and 2.283 minutes, respectively. The peak of target compound was visibly not present in the control chromatogram at the stated retention times (Fig. 3).

### 4. DISCUSSION

Results of the study indicated the presence of the dichlorvos six months post application in the treated cowpea grain. The detection of dichlorvos in the treated grains could be attributed to the inappropriate application such as over dosing of the said pesticide done indiscriminately to achieve set goals [14]. Most farmers (especially grain marketers) exposed cowpea grains to dangerous chemicals especially organophosphates which are normally higher or above normal approved doses by the governments/or international organizations such as FAO, WHO etc [14]. This improper use of pesticides may lead to the emergence of resistant pest populations, which would be insensitive to normal pesticide dosages [14, 15 and 16].

A study in Zimbabwe have indicated that illiteracy, poor or bad attitudes of farmers on dangers of pesticides contributed to higher levels of dichlorvos in vegetables in Zimbabwe [17] and such could be a similar trend happening at Dawanau market in Kano State. Higher dosages






Fig. 1. Chromatogram of the dichlorvos protected cowpea grains



**Fig. 2. Chromatogram of a commercially available brand of dichlorvos used as standard**



**Fig. 3. Chromatogram of the unprotected cowpea grains used as control**

of Dichlorvos are being used irrespective of the recommended dosage in grains storage in the sub Saharan Africa especially in Dawanau grains Market each year as both farmers and traders alike await better prices. It is equally a common practice that calendar application of such pesticides are widely practiced by both farmers and traders, this particular practice is capable of increasing pesticide residue accumulation in such stored grains [2]. The grain traders/merchants are directly mixing grains with

higher dosages of chemical insecticides in order to have maximum and long term protection against insect pest damage. The detected levels of dichlorvos on the treated cowpea grain could be due to shortening of the recommended withdrawal period, which is the minimum time one must wait between pesticide application and usage by the ultimate consumers of any insecticide treated commodity. Organophosphorous pesticides are widely applied in agriculture because their degradation

elsewhere is known to be rapid [18]. However, in this study Dichlorvos is said to persist for six months after application. This could be attributed to the higher dosage of the pesticide applied.

In Nigeria, especially in the northern part, dichlorvos is traded under different names such as Nuvan, Sniper or "Pia-pia" or "Otah Pia-pia" (a local name) and is handled and used as a household insecticide indiscriminately. Moreover, excessive or higher dosages of Dichlorvos applied by traders and farmers in Dawanau market have probably caused the persistence and possibly pest resistance in stored grains protected using dichlorvos. This has been reported by several researchers including [19] that the stored product insects pests were found to be resistant against different insecticides including the dichlorvos, cyclodienes, chlorpyrifos, cyanophos, carbamates, carbaryl, cypermethrin, dichlorodiphenyltrichloroethane, deltamethrin, diazinon, ethylene bromide, ethyl formate organophosphates, permethrin, pyrethrins, and propoxur. More so, [20] reported that resistance to pesticides used to protect grain and other stored food stuffs is widespread and involves all groups of pesticides and most of the important pests. The residue of dichlorvos found in the stored cowpea grains may lead to various health issues that are unknown by the grain marketers or the final consumers. This might expose the final consumer to different diseases such as impaired memory and concentration, disorientation, severe depressions, irritability, confusion, headache, speech difficulties, delayed reaction times, nightmares, sleepwalking and drowsiness or insomnia. An influenza-like condition with headache, nausea, weakness, loss of appetite, and malaise has also been reported [21,14]. In a similar study by [22], results clearly indicated that inhalation of chlorpyrifos, dichlorvos and alphacypermethrin for eight weeks caused a slight decrease in the average body weight gain in test mice.

## 5. CONCLUSION

The present investigation clearly indicated the presence of dichlorvos (i.e. an organophosphate insecticide) in cowpea grains six months after application and this situation highlights among others, the dangers and/or disadvantages associated with the use of synthetic chemicals used for the protection of stored cowpea grains against insect pests especially by the personnel lacking requisite knowledge on the appropriate ways of handling/application of such products.

Owing to the serious implications of the inappropriate use of synthetic chemicals on humans and environment there is therefore, the need for the regulatory authorities to target places like Dawanau market with campaign strategies on best practices in the use of such chemicals on agricultural commodities. More so, further research or researches should be conducted especially on the maximum residue limits (MRL) and its comparison with the Codex Alimentarius as well as the extension in the storage period.

## ACKNOWLEDGEMENTS

We sincerely thank the Vice-Chancellor and the entire management team of Bayero University, Kano for funding the research work.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Adedire CO. Biology, ecology and control of insect pests of stored cereal grains. In: *Pests of Stored Cereals and Pulses in Nigeria-biology, ecology and control* (ed) Ofuya, T.I. and Lale, N.E.S. Nigeria: Dave Collins Publications. 2001;174.
2. Dahiru B, Abdullahi G, Bukar N. Pesticides use among grain merchants in Mubi; 2014.
3. Grain markets of Adamawa State, Nigeria. *Agroresearch*. 2014;14(1):1–13.
4. Xiaosong G, Weston PA. Oviposition and feeding deterrents from Chinese prickly ash against Angoumois grain moth (Lepidoptera: Gelechiidae). *Journal of Economic Entomology*. 1995;88:1771-1775.
5. Aaron CK. Organophosphates and carbamates. In: Ford MD, Delaney KA, Ling LJ, Erickson T, (ed). *Clinical toxicology*. WB Saunders Company; Philadelphia: 2001;819–828.
6. Kennedy IR. Biodegradation and plant protection from herbicide 2, 4-D by plant Microbial Association in Cotton. 1998;30-41.
7. Okori P, Rubaihayo PR, Adipala E, Fahlesion J, Dixelium C. Population studies of fungal plant pathogen; perspectives for control with specific

- reference to grey leaf spot. Africa Crop Science Journal. 2004;12(4):243-249.
8. Czarniewska E, Kasprzyk A, Ziemnicki K. Effect of paraquat and metoxychlor on antioxidant enzymes in frog *Rana esculenta*. L. liver. Biological Letters. 2003; 2:125-133.
  9. Jean S, Ngassoum MB, Tchatchueng JB, Ashutosh KS, Laxman PS. Contamination of cowpea and by-products by organophosphorous pesticide residues in Ngaoundere markets: Dietary risk estimation and degradation study. African Journal of Food Science. 2013;7(5):92–102.  
DOI: 10.5897/AJFS2013.1013
  10. Jeyaratnam J. Acute pesticide poisoning: A major global health problem. World Health Stat Q. 1990;43(3):139-44.
  11. Nguengang P, Parrot L, Lejoly-Joiris VJ. Yaounde Dregs exploitation: Production system, traditional ability and urban and peri-urban agricultural potentialities for local development. In: Agriculture and urban development in West and Central Africa. L. Parrot and R. Kahane, eds. Montpellier, CIRAD, France. 2005;40.
  12. Lotti M. Clinical toxicology of anticholinesterase agents in humans. In: Krieger R, (ed). Handbook of pesticide toxicology. Agents. 2nd ed. Academic Press; San Diego. 2001;2:1043–1085.
  13. Akoto O, Andoh H, Darko G, Eshun K, Osei-Fosu P. Health risk assessment of pesticides residue in maize and cowpea from Ejura, Ghana.  
DOI:10.1016/j.chemosphere.2013. 02.057 PubMed.
  14. Chedi BAZ, Aliyu M. Effect and management of acute dichlorvos poisoning in wistar rats. Bayero Journal of Pure and Applied Sciences. 2010;3(2): 1–3.
  15. Davies MS, Boniface M, Gibson S. Determination of dichlorvos residue levels in vegetables sold in Lusaka, Zambia. PanAfrican Medical Journal. 2016; 23:113.8211
  16. Bardin M, Ajouz S, Comby M, Lopez-Ferber M, Graillot B, Siegwart M, Nicot PC. Is the efficacy of biological control against plant diseases likely to be more durable than that of chemical pesticides? Front Plant Sci. 2015, PubMed | Google Scholar
  17. Li SG. Drug resistance evolution of dichlorvos-resistant and cypermethrin-resistant strains of *Culex pipiens pallens*. Zhongguo ji sheng chong xue yu ji sheng chong bing za zhi. Chinese Journal of Parasitology & Parasitic Diseases. 2009; 27(4):311-312.
  18. Kavi KS, Reetha D. Efficiency of Bacterial Isolates in the Degradation of Malathion and Parathion. Int. J. Pharm. Biol. Arch. 2012;3(3):659–665.
  19. Rajashekar Y, Nandagopal B, Thimmappa S. Botanicals as Grain Protectants (A review). Psyche. 2012;1–13.
  20. Champ BR. Occurrence of resistance to pesticides in grain storage pests, In Pesticides and Humid Tropical Grain Storage System, B. R. Champ and E. Highly, Eds., of Proceedings of an International Seminar, Manila 1985. Canberra ACIAR Proceedings. 1985;14: 225–229.
  21. Clark RF. Goldfrank's toxicological emergencies. 7th ed. McGraw-Hill Professional; New York: Insecticides: organic phosphorus compounds and carbamates. 2002;1346–1360.
  22. Kemabonta KA, Akinhanmi FO. Toxicological effects of chlorpyrifos, dichlorvos and alpha cypermethrin on adult albino mice, *Mus musculus*. Production Agriculture and Technology. 2013;9(2):1–17.

© 2017 Yusuf et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://sciencedomain.org/review-history/22570>